


Mitigating Necrotic Enteritis through Natural Alternatives in Antibiotic-Free Production Systems



by EW Nutrition USA, Inc.

In the poultry industry, Necrotic Enteritis is of great interest due to the potential detrimental growth effects it may have in a flock, even at subclinical levels⁵⁰. Coccidiostats and antibiotics have been used for a long time to get the disease-causing bacterium *Clostridium perfringens* under control, but with increasing antimicrobial resistance, alternative approaches are required. This article aims to give an overview of the disease and the measures against it.



Clostridium perfringens – a ubiquitous, highly resilient bacterium

Clostridium perfringens is a Gram-positive, spore-forming, anaerobic, rod-shaped bacterium⁵⁰. This encapsulated, non-motile microorganism is fastidious in growth requirements⁵⁹. Most often, complex media like cooked meat or thioglycolate broth are used as enrichment³⁰.

It was Welch and Nuttall who first identified *C. perfringens* in 1892 as *Bacillus aerogenes capsulatus*¹⁸. In Great Britain, the bacterium was commonly known as *C. welchii* and sometimes called Frankel's bacillus in Germany until designated *C. perfringens* by Bergey¹³.

Clostridium perfringens is the causal microorganism for Necrotic Enteritis (NE)¹⁴. In humans, it is one of the most common causes of foodborne illness²⁰. The Centers for Disease Control and Prevention (CDC, 2012) estimates that nearly one million people are affected every year, making *C. perfringens* the third most frequent source of domestically acquired foodborne illness after Norovirus and *Salmonella*.

Clostridium perfringens can be found everywhere

Clostridium perfringens is found in soil, water, and other organic materials. As far as poultry facilities, *C. perfringens* has been isolated from litter, dust, walls, floors, fans, transportation coops, feeders, and feed⁸⁹.

Additionally, *C. perfringens* is found in the GI tract of broiler chickens, humans, and other mammals⁴⁷. When intestinal samples of broiler chickens were analyzed for *C. perfringens*, 75-95 % tested positive²⁴.

Drew and co-workers¹⁰ determined that *C. perfringens* is usually found at $\sim 10^4$ colony-forming units (CFU)/g of broiler digesta. These results agree with Jia et al.²⁶, who stated that *C. perfringens* is present at low levels in healthy poultry. In humans, investigations in different parts of the world showed a prevalence of *Clostridium perfringens* between 57-94%³².

Different types of *Clostridium perfringens* with different toxins

There are five types (A-E) of *C. perfringens*, which can be identified through their toxin production (see table 1). All strains produce alpha-toxin. Furthermore, *Clostridium perfringens* has been described to produce eight other toxins, three (delta, theta, kappa) can be lethal, but these are seldom involved in disease origin³⁷.

Table 1. Different types of *Clostridium perfringens*

		<i>C. perfringens</i> Type				
		A	B	C	D	E
Toxins	Alpha	x	x	x	x	x
	Beta		x	x		
	Epsilon		x		x	
	Iota					x
	Enterotoxin	x				
Diseases/animals ¹⁸	Food-born disease/humans NE/fowl	Dysentery/lambs enterotoxaemia/ sheep, goats, guinea pigs	Food-born disease/humans NE/fowl	Enterotoxaemia/ sheep Pulpy kidney disease/lambs	Enterotoxaemia/ calves Dysentery/sheep, guinea pigs, rabbits	

High resilience gives an advantage against competitors

Since *Clostridium perfringens* is a spore-forming bacterium, it is very resilient to high temperatures, slight pH variations, and toxic chemicals^{43, 7}.

Labbe et al.³⁰ established that *C. perfringens* can reproduce at temperatures between 15-50 °C. Hence, proper refrigeration temperatures (below 10 °C) can be an effective means of control. The optimum range is between 37-47 °C, and at these temperatures, the mean generation time - the time required for the bacterial count to double - is approximately 10-12 minutes⁴¹. These short generation times allow the bacteria to outcompete other microorganisms that may need similar resources in a certain environment.

The optimum pH range of *Clostridium perfringens* is between 5.5-7.0²². However, it can grow at a pH as low as 5 and as high as 9. In live broiler chickens, the pH in the small intestine has been determined to be between 6.00-7.78.

Necrotic enteritis in poultry

The disease necrotic enteritis was first described by Parish^{45, 46} in cockerels in England. Some of the symptoms include depression, reluctance to move, ruffled feathers, somnolence, diarrhea, loss of appetite, and anorexia²¹. Mortality ranges from 0-50%⁶ have been reported in infected flocks. Since then, virtually every area that raises poultry has reported signs of necrotic enteritis.

Clostridium perfringens – How NE unravels

As already mentioned, 10^4 colony-forming units (CFU)/g of broiler digesta¹⁰ are normal and can be found in healthy birds. *C. perfringens* becomes problematic when counts reach 10^7 - 10^8 CFU/g⁶.

Necrotic enteritis is caused by types A and C of *Clostridium perfringens*, but normally, predisposing factors “set the stage”^{24, 48}. This could be seen in an investigation where they wanted to create a model to reproduce NE in a laboratory setting. Researchers realized that inoculation of *C. perfringens* alone did not cause the disease found in the field⁴⁸. Therefore, it was assessed that certain cofactors must play a significant role in the pathogenicity of *C. perfringens*. Williams⁵⁷ reviewed concurrent infections of coccidiosis and necrotic enteritis in chickens (Figure 1). The copious interactions of these diseases with predisposing factors, control methods, sources of infection, and disease form is a testament to the complexity of this poultry industry matter.

Coccidiosis creates access

Shane *et al.*⁵³ noted that several authors had considered coccidiosis to be a predisposing factor for NE. They proceeded to describe the pathogenesis of *Eimeria acervulina*, one of the protozoa responsible for coccidiosis in poultry. When the oocysts are ingested, they quickly attach to the intestinal wall causing lesions where the protozoa reproduce numerous times. These are the lesions to which *C. perfringens* attaches.

What happens in the animal?

Long *et al.*³³ proposed the pathogenesis for NE: First, epithelial cells are vacuolated, and the epithelium lifts off the lamina propria, which is congested and edematous. These lesions can be caused by a combination of factors like toxin production and/or, as just mentioned, coccidiosis. *Clostridium perfringens* cells attach to the lamina propria, where they thrive. The tissue becomes necrotic as large numbers of heterophils, a type of phagocyte, flood the foci (sites of lesions).

A combination of disease-inducing factors such as bacteria proliferation, heterophil lysis, and villus' necrosis seem to develop quickly. The inflammation zone then becomes riddled with mononuclear cells, cells containing lymphocytes, antigen-presenting cells, and eosinophilic-staining (proteinaceous) amorphous material. This necrotizing process moves from the tip of the villi to the crypt.

Chronic version

In chronic cases, villi may be found to have multiple cysts from recurrent necrosis. In birds that overcome the disease, injured epithelial cells are replaced by newly formed reticular structures. These new cells travel from the crypt to the tip of the villi and replace the old, damaged cells. The result is a short, flat villus with a reduced surface area for nutrient absorption^{44, 45, 34}. These morphologically altered villi are the

necrotic lesions found in the field and some *C. perfringens* challenge trials (Figure 2).

Acute form

The acute form of NE results in enlarged lesions along the gut wall, and the epithelium becomes eroded and detached; consequently, a diphtheritic membrane is formed. This yellow, green, or brownish pseudo-membrane is called the “Turkish towel,” which describes the appearance of the friable, gas-filled, foul-smelling GI tract⁵⁷.

Subclinical form

Poultry producers are not only concerned with the acute form of NE. Recent studies have shown that the disease’s subclinical form can be as detrimental as the acute illness¹⁹. Lovland and co-workers³⁵ stated that this symptomless disease is often overlooked at the farm, and the effects are only noticed at the processing facility.

Subclinical NE (SNE) can cause cholangiohepatitis, a condition where the liver is enlarged with pale reticular patterns and sometimes small, pale foci. In the United Kingdom, it was estimated that 4% of broiler carcasses and 12% of livers are condemned at processing plants due to clostridial infection; thereby, reducing profit³⁶. Moreover, sparse lesions that may be found in a case of SNE may be enough to hinder growth performance; thus, resulting in an underproductive flock³⁹.

Feeding Against Necrotic Enteritis

It has been reported that diet formulation has the greatest impact on the prevalence of *C. perfringens* in chicken GI tracts⁶¹. The poultry industry formulates diets on a least-cost basis, which may become problematic if nutritionists do not take into consideration the pathological consequences that some ingredients may have in the GI tracts of chickens. Every feed ingredient has a specific purpose in the diet. For instance, cereal grains are fed for their energy concentration as well as fiber. Also, some grain and animal/plant meals are used for their protein content. Since these ingredients are obtained from different sources, they are highly variable in macro and micronutrients¹.

The diet provides the conditions for proliferation

There are multiple elements that affect the proliferation of *C. perfringens* in chicken intestines, one of the most critical factors being diet formulation^{5, 36}. Some feed ingredients have been found to exacerbate the numbers of *C. perfringens* in chickens’ gastrointestinal tract. Diets formulated with wheat increased NE intestinal lesion scores compared to broiler chickens fed a corn-based diet⁴. In another study, Drew *et al.*¹⁰ investigated the effects of different protein sources on the intestinal populations of *C. perfringens* in broiler chickens. Diets were formulated to contain 230, 315, and 400 g/kg of fishmeal or soy protein concentrate (SPC). The numbers of *C. perfringens* in the ileum and ceca increased when the amount of protein increased from 230 to 400 g/kg.

Type of grain influences the occurrence of *Clostridium perfringens*

Authors have studied the effects of grain inclusion on gut microbiota, and it is well established that small cereal grains such as barley, rye, and wheat tend to increase the prevalence of *C. perfringens* in the GI tract. Shakouri *et al.*⁵² investigated the influence of barley, sorghum, wheat, and corn on counts of *C. perfringens* in the different intestinal segments. Corn and wheat had the lowest *C. perfringens* counts, followed by sorghum, while barley yielded the highest counts. These findings agree with Riddell and Kong⁵¹.

Other researchers have concluded that the increase in gut viscosity and increased chyme transit time elicit the overgrowth of *C. perfringens* in the intestines²⁸. Grains like wheat and barley contain high amounts of non-starch polysaccharides (NSP), which increase viscosity²⁶. Furthermore, it has been alleged that, since these grains are high in NSP, the bird cannot absorb nutrients as efficiently, thereby leaving them for microbes like *C. perfringens* to consume³¹.

Enzymes improve nutrient availability in the presence of *C. perfringens*

Shakori *et al.*⁵² and Jia *et al.*²⁶ also studied the impact of several diets with the inclusion of a blend of carbohydrases such as glucanase and xylanase. Their findings suggested that enzyme addition did not affect counts of *C. perfringens* in the different intestinal sections. However, they did find an improvement in growth performance. They stated that enzymes improved chyme viscosity by degrading the encapsulation of nutrients in diets.

For this reason, researchers have investigated the use of enzymes in wheat and barley-based diets on the incidence of *C. perfringens* in chicken intestines. Jackson *et al.*²⁵ studied the effect of beta-mannanase addition on flocks infected with *Eimeria* spp. and *C. perfringens*. They found that feeding this enzyme significantly reduced the impact of *C. perfringens* on the performance of infected flocks as well as intestinal lesion scores. Moreover, the authors explained that this might be due to beta-mannanase crossing the intestinal wall to provoke an immune response. They determined that this enzyme tended to ameliorate the symptoms of necrotic enteritis, but not significantly.

MOS may have a positive impact on immunity

Hofacre *et al.*²³ found similar results when birds were fed mannan-oligosaccharides. A marked effect was only found when mannan-oligosaccharides were included along with lactic acid-producing, competitive exclusion products (probiotics).

The feed form is decisive

Feed form has also been investigated on the incidence of *C. perfringens*. When birds were fed whole wheat compared to ground, researchers found reduced counts of *C. perfringens* in the gut². These results can be extrapolated to the findings of Engberg *et al.*¹¹. They found that when birds were fed coarse versus fine mash or pellets, *C. perfringens* counts were consistently higher in flocks fed mash diets. These authors concluded that feeding pellets or whole grains increases gizzard activity, which consequently triggers

hydrochloric acid production and decreases pH in the GI tract. This drop in pH of approximately 0.5 units may be responsible for decreased *C. perfringens* counts.

Mind the protein source

Another well-established fact is that the *C. perfringens* population can be affected by the type of the protein source and the inclusion rates.

Potato is worse than fish

Palliyeguru *et al.* ⁴² studied the inclusion of protein concentrates (potato, fish, and soy) on subclinical NE. They determined that the potato-containing diet resulted in the highest incidence of *C. perfringens* in the gut, followed by fish and soy. Also, the potato-containing diet had the highest activity of trypsin inhibitors and lowest lipid content. Increased trypsin inhibition does not allow for the inactivation of alpha and beta toxins produced by *C. perfringens*, resulting in increased intestinal wall lesions.

Fish is worse than soy due to the amino acid composition

Drew *et al.* ¹⁰ formulated diets containing fishmeal or a soy protein concentrate at different levels. Feeding dietary fishmeal resulted in a higher incidence of *C. perfringens* as compared to the soy protein diet. Furthermore, with increasing levels of soy and fishmeal diets, counts of *C. perfringens* increased as well. A notable difference in fishmeal protein concentrate compared to the soy protein concentrate was the amino acid ratio in this experiment; the methionine and glycine ratios were 1.3 times greater in fishmeal diets.

Muhammed *et al.* ⁴⁰ determined that methionine was required for *C. perfringens* sporulation. This may be of interest to nutritionists since some authors have estimated that 10-20 % of synthetic amino acids are not absorbed and reach the lower intestinal tract, i.e., ceca; thereby, aiding in the proliferation of *C. perfringens*.

Fat source – animal fat is critical

The effects of fat sources on *C. perfringens* population remain largely unknown. Knarreborg *et al.* ²⁹ studied the bacterial microflora in chicken intestines after feeding different dietary fats (soy oil and a tallow and lard mix) in rations containing antibiotic growth promoters (AGP). When soy oil was fed, *C. perfringens* counts were significantly lower than diets containing animal fats. The authors stated that, since plant oils contain higher amounts of unsaturated fatty acids, the chyme in birds fed oil diets would have decreased viscosity, decreasing transit time. Furthermore, an additive effect was found when soy oil was provided along with AGP, which may be due to facilitated antibiotic dispersion caused by the oil's lipophilic properties. Knarreborg *et al.* (2002) investigated the effects of fat sources on *C. perfringens*. They found that total anaerobic counts increased with animal fat addition. However, zinc bacitracin was included in their diets, specifically targeting Gram-positive microorganisms like *C. perfringens*; thus, potentially biasing their results.

Antibiotics and coccidiostats in the diet –

helpful, but finite

Antibiotics and coccidiostats have been commonly included in poultry diets since the mid-1940s and 1950s^{61, 58}.

Prescott *et al.*⁴⁹ studied the inclusion of zinc bacitracin to prevent necrotic enteritis and concluded that it successfully controlled the *C. perfringens* challenge. Flocks in the antibiotic treatments were able to overcome disease and perform similarly to unchallenged birds. Multiple authors have replicated these results using different antibiotics such as virginiamycin and salinomycin^{17, 3, 11}.

Improvements in flock performance with the inclusion of antibiotics and coccidiostats are well understood and omnipresent in the literature. However, the potential loss of subtherapeutic antibiotic usage in livestock in the United States due to increasing concerns over [antimicrobial resistance](#) and consumer demands makes research of viable alternatives to these compounds paramount.

So, what are your alternatives?

A lot of different approaches are possible. In general, these measures should act against *Clostridium perfringens* while supporting gut health.

Tested substances without the desired effects

Lastly, multiple options have been studied to control *C. perfringens* in poultry. Some researchers have studied the inclusion of complex carbohydrates and fibers like pine shavings, guar gum, and pectin with limited success^{4, 31}. Another popular alternative is the use of competitive exclusion-based products such as prebiotics and probiotics^{27, 16}. Still, these products failed to yield consistent results.

Other options that have been investigated are the addition of lactose and organic acids^{54, 38}. Potassium diformate did not produce lowered counts of *C. perfringens*. Lactose reduced *C. perfringens* counts but resulted in undesirable ceca characteristics including, enlargement and increased fermentation⁵⁴.

Essential oils alone or in combination may be a solution

Mitsch and coworkers³⁹ investigated the efficacy of two blends of essential oils with positive effects on the reduction of *C. perfringens* from the gut and feces of broilers. Gaucher and coworkers¹⁵ compared growth performance and gut health of broilers fed a conventional (anticoccidials and AGPs) vs. ABF (Coccidiosis vaccine and essential oil blends) diet. They established that livability, age at slaughter, and percentage of condemnation did not change with diet type. However, average daily weight gain and FCR were negatively affected. Furthermore, NE was more prevalent in ABF flocks. Still, many authors agree that a multifactorial approach is necessary if antibiotics should be completely replaced by these strategies³⁶.

A contemporary study by Wati *et al.*⁵⁶ aimed to compare AGPs to a commercial blend of essential oils fed to broilers. Authors found that chickens fed essential oils had body weights and FCRs that were statistically similar to the AGP treatment. Moreover, both AGP and essential oil treatments had statistically lower counts of *Salmonella* and *E. coli* after an oral challenge than the control group.




Figure 2. Necrotic Enteritis lesions in chicken intestines

Yellowish necrotic lesions in three intestinal samples. Intestines A and C show a few marked lesions. Intestine B shows clusters of lesions typical of the “Turkish towel” syndrome. (Source: <http://www.mdpi.com/2072-6651/2/7/1913/htm>. Accessed: January 14, 2021).

References

1. Bedford, M.R. 1996. Interaction Between Ingested Feed and the Digestive System in Poultry. *Applied Poultry Science* 5:86-95.
2. Bjerrum, L., K. Pedersen, and R. M. Engberg. 2005. The Influence of Whole Wheat Feeding on Salmonella Infection and Gut Flora Composition in Broilers. *Avian Disease* 49:9-15.
3. Bolder, N. M., J. A. Wagenaar, F. F. Putirulan, K. T. Veldman, and M. Sommer. 1999. The Effect of Flavophospholipol (Flavomycin[®]) and Salinomycin Sodium (Sacox[®]) on the Excretion of *Clostridium perfringens*, *Salmonella enteritidis*, and *Campylobacter jejuni* in Broilers After Experimental Infection. *Poultry Science* 78:1681-1689.
4. Branton, S. L., B. D. Lott, J. W. Deaton, W. R. Maslin, F. W. Austin, L. M. Pote, R. W. Keirs, M. A. Latour, and E. J. Day. 1997. The Effect of Added Complex Carbohydrates or Added Dietary Fiber on Necrotic Enteritis Lesions in Broiler Chickens. *Poultry Science* 76:24-28.
5. Choct, M. 2009. Managing Gut Health through Nutrition. *British Poultry Science* 50:9-15.
6. Cooper, K., and J. G. Songer. 2009. Necrotic Enteritis in Chickens: A Paradigm of Enteric Infection by *Clostridium perfringens* Type A. *Veterinary anaerobes and diseases* 15:55-60.
7. Craven, S. E., N. J. Stern, N. A. Cox, J. S. Bailey, and M. Berrang. 1999. Cecal Carriage of *Clostridium perfringens* in Broiler Chickens Given Mucosal Starter Culture[™]. *Avian Diseases* 43:484-490.
8. Craven, S. E., N. A. Cox, N. J. Stern, and J. M. Mauldin. 2001a. Prevalence of *Clostridium perfringens* in Commercial Broiler Hatcheries. *Avian Diseases* 45:1050-1053.
9. Craven, S. E., N. J. Stern, J. S. Bailey, and N. A. Cox. 2001b. Incidence of *Clostridium perfringens* in Broiler Chickens and Their Environment during Production and Processing. *Avian Diseases* 45:887-896.
10. Drew, M. D., N. A. Syed, B. G. Goldade, B. Laarveld, and A. G. Van Kessel. 2004. Effects of Dietary Protein Source and Level on Intestinal Populations of *Clostridium perfringens* in Broiler Chickens. *Poultry Science* 83:414-420.
11. Engberg, R. M., M. S. Hedemann, Leser, T.D., and Jensen, B.B. 2000. Effect of Bacitracin and

- Salinomycin on Intestinal Microflora and Performance of Broilers. Poultry Science 79: 1311-1319.
12. Engberg, R. M., M. S. Hedemann, and Jensen, B.B. 2002. The Influence of Grinding and Pelleting of Feed on the Microbial Composition and Activity in the Digestive Tract of Broiler Chickens. British Poultry Science 44:569-579.
 13. Freeman, B.A. 1979. *Burrows Textbook of Microbiology*. W.B. Saunders Company, Philadelphia, Pennsylvania, USA.
 14. Fukata, T., Y. Hadate, E. Baba, T. Uemura, and A. Arakawa. 1988. Influence of *Clostridium perfringens* and its Toxin in Germ-free Chickens. Research in Veterinary Science 44:68-70.
 15. Gaucher, M.L., Quessy, S., Letellier, A., Arsenault, J., and M. Boulianne. 2015. Impact of a drug-free program on broiler chicken growth performances, gut health, *Clostridium perfringens* and *Campylobacter jejuni* occurrences at the farm level. Poultry Science 94: 1791-1801.
 16. Geier, M. S., L. L. Mikkelsen, V. A. Torok, G. E. Allison, G. C. Olmood, M. Boulianne, R. J. Hughes, and M. Choct. 2010. Comparison of Alternatives to In-feed Antimicrobials for the Prevention of Clinical Necrotic Enteritis. Journal of Applied Microbiology 109:1329-1338.
 17. George, B. A., C. L. Quarles, and D. J. Fagerberg. 1982. Virginiamycin Effects on Controlling Necrotic Enteritis Infection in Chickens. Poultry Science 61:447-450.
 18. Hatheway, C. L. 1990. Toxigenic Clostridia. Clinical Microbiology Reviews 3:66-98.
 19. Heier, B. T., A. Lovland, K. B. Soleim, M. Kaldhusdal, and J. Jarp. 2001. A Field Study of Naturally Occurring Specific Antibodies against *Clostridium perfringens* Alpha Toxin in Norwegian Broiler Flocks. Avian Diseases 45:724-732.
 20. Heikinheimo, A., M. Lindstrom, and H. Korkeala. 2004. Enumeration and Isolation of *cpe*-Positive *Clostridium perfringens* Spores from Feces. Journal of Clinical Microbiology 42:3992-3997.
 21. Helmboldt, C. F., and E. S. Bryant. 1971. The Pathology of Necrotic Enteritis in Domestic Fowl. Avian Diseases 15:775-780.
 22. Hickey, C.S., and Johnson, M.G. 1981. Effects of pH Shifts, Bile Salts, and Glucose on Sporulation of *Clostridium perfringens* NTCT 8798. Applied and Environmental Microbiology 41:124-129.
 23. Hofacre, C. L., T. Beacorn, S. Collett, and G. Mathis. 2003. Using Competitive Exclusion, Mannan-Oligosaccharide and Other Intestinal Products to Control Necrotic Enteritis. Journal of Applied Poultry Research 12:60-64.
 24. Immerseel, F. V., J. De Buck, F. Pasmans, G. Huyghebaert, F. Haesebrouck, and R. Ducatelle. 2004. *Clostridium perfringens* in Poultry: an Emerging Threat for Animal and Public Health. Avian Pathology 33:537-549.
 25. Jackson, M. E., D. M. Anderson, H. Y. Hsiao, G. Mathis, and D. W. Fodge. 2003. Beneficial Effect of B-Mannanase Feed Enzyme on Performance of Chicks Challenged with *Eimeria* and *Clostridium perfringens*. Avian Diseases 47:759-763.
 26. Jia, W., B. A. Slominski, H. L. Bruce, G. Blank, G. Crow, and O. Jones. 2009. Effects of Diet Type and Enzyme Addition on Growth Performance and Gut Health of Broiler Chickens During Subclinical *Clostridium perfringens* Poultry Science 88:132-140.
 27. Kaldhusdal, M., C. Schneitz, M. Hofshagen, and E. Skjerve. 2001. Reduced Incidence of *Clostridium perfringens*-Associated Lesions and Improved Performance in Broiler Chickens Treated with Normal Intestinal Bacteria from Adult Fowl. Avian Diseases 45:149-156.
 28. Klasing, K. C. 1998. Nutritional Modulation of Resistance to Infectious Diseases. Poultry Science 77:1119-1125.
 29. Knarreborg, A., M. A. Simon, R. M. Engberg, B. B. Jensen, and G. W. Tannock. 2002. Effects of Dietary Fat Source and Subtherapeutic Levels of Antibiotic on the Bacterial Community in the Ileum of Broiler Chickens at Various Ages. Applied and Environmental Microbiology 68:5918-5924.
 30. Labbe, R. G. 1991. *Clostridium perfringens*. Journal of the Association of Official Analytical Chemists 74:711-714.
 31. Langhout, D. J., J. B. Schutte, P. Van Leeuwen, J. Wiebenga, and S. Tamminga. 1999. Effect of Dietary High- and Low-methylated Citrus Pectin on the Activity of the Ileal Microflora and Morphology of the Small Intestinal Wall of Broiler Chicks. British Poultry Science 40:340-347.
 32. Lindstrom, M., A. Heikinheimo, P. Lahti, and H. Korkeala. 2011. Novel Insights into the Epidemiology of *Clostridium perfringens* Type A Food Poisoning. Food Microbiology 28:192-198.
 33. Long, J.R., Pettit, J.R., and Barnum, D.A. 1974. Necrotic Enteritis in Broiler Chickens II. Pathology and Proposed Pathogenesis. Canadian Journal of Comparative Medicine 38: 467-474.
 34. Long, J. R., and R. B. Truscott. 1976. Necrotic Enteritis in Broiler Chickens III. Reproduction of the Disease. Canadian Journal of Comparative Medicine 40:53-59.
 35. Lovland, A., and M. Kaldhusdal. 1999. Liver Lesions Seen at Slaughter as an Indicator of Necrotic Enteritis in Broiler Flocks. FEMS Immunology and Medical Microbiology 24:345-351.
 36. McDevitt, R. M., J. D. Brooker, T. Acamovic, and N. H. C. Sparks. 2006. Necrotic Enteritis; A Continuing Challenge for the Poultry Industry. World's Poultry Science Journal 62:221-247.

37. McDonel, J. L. 1986. *Clostridium perfringens* Toxins (type A, B, C, D, E). *Pharmacology and Therapeutics* 10:617-655.
38. Mikkelsen, L. L., J. K. Vidanarachchi, G. C. Olnood, Y. M. Bao, P. H. Selle, and M. Choct. 2009. Effect of Potassium Diformate on Growth Performance and Gut Microbiota in Broiler Chickens Challenged with Necrotic Enteritis. *British Poultry Science* 50:66-75.
39. Mitsch, P., K. Zitterl-Eglseer, B. Kohler, C. Gabler, R. Losa, and I. Zimpf. 2004. The Effect of Two Different Blends of Essential Oil Components on the Proliferation of *Clostridium perfringens* in the Intestines of Broiler Chickens. *Poultry Science* 83:669-675.
40. Muhammed, S. I., S. M. Morrison, and W. L. Boyd. 1975. Nutritional Requirements for Growth and Sporulation of *Clostridium perfringens*. *Journal of Applied Bacteriology* 38:245-253.
41. Murray, P. R., K. S. Rosenthal, and M. A. Pfaller. 2009. *Medical Microbiology*. 6th ed. Elsevier Health Sciences, Philadelphia, PA, USA.
42. Palliyeguru, M. W. C. D., S. P. Rose, and A. M. Mackenzie. 2010. Effect of Dietary Protein Concentrates on the Incidence of Subclinical Necrotic Enteritis and Growth Performance of Broiler Chickens. *Poultry Science* 89:34-43.
43. Paredes-Sabja, D., Torres, J.A., Setlow, P., and Sarker, M.R. 2008. *Clostridium perfringens* Spore Germination: Characterization of Germinants and their Receptors. *Journal of Bacteriology* 190:1190-1201.
44. Parish, W. E. 1961. Necrotic Enteritis in the Fowl (*Gallus Gallus Domesticus*). I. Histopathology of the Disease and Isolation of a Strain of *Clostridium welchii*. *Journal of Comparative Pathology* 71:377-393.
45. Parish, W. E. 1961. Necrotic Enteritis in the Fowl. II. Examination of the Causal *Clostridium welchii*. *Journal of Comparative Pathology* 71:394-404.
46. Parish, W. E. 1961. Necrotic Enteritis in the Fowl. III. The Experimental Disease. *Journal of Comparative Pathology* 71:405-414.
47. Pedersen, K., L. Bjerrum, B. Nauerby, and M. Madsen. 2003. Experimental Infections with Rifampicin-resistant *Clostridium perfringens* Strains in Broiler Chickens Using Isolator Facilities. *Avian Pathology* 32:403-411.
48. Pedersen, K., L. Bjerrum, O. Heuer, D. Wong, and B. Nauerby. 2007. Reproducible Infection Model for *Clostridium perfringens* in Broiler Chickens. *Avian Diseases* 52:34-39.
49. Prescott, J. F., R. Sivendra, and D. A. Barnum. 1978. The Use of Bacitracin in the Prevention and Treatment of Experimentally-induced Necrotic Enteritis in the Chicken. *Canadian Veterinary Journal* 19:181-183.
50. Rehman, H., W. A. Awad, I. Lindner, M. Hess, and J. Zentek. 2006. *Clostridium perfringens* Alpha Toxin Affects Electrophysiological Properties of Isolated Jejunal Mucosa of Laying Hens. *Poultry Science* 85:1298-1302.
51. Riddell, C., and X. Kong. 1992. The Influence of Diet on Necrotic Enteritis. *Avian Diseases* 36:499-503.
52. Shakouri, M. D., P. A. Iji, L. L. Mikkelsen, and A. J. Cowieson. 2008. Intestinal Function and Gut Microflora of Broiler Chickens as Influenced by Cereal Grains and Microbial Enzyme Supplementation. *Journal of Animal Physiology and Animal Nutrition* 93:647-658.
53. Shane, S. M., J. E. Gyimah, K. S. Harrington, and T. G. Snider. 1985. Etiology and Pathogenesis of Necrotic Enteritis. *Veterinary Research Communications* 9:269-287.
54. Takeda, T., T. Fukata, T. Miyamoto, K. Sasai, E. Baba, and A. Arakawa. 1995. The Effects of Dietary Lactose and Rye on Cecal Colonization of *Clostridium perfringens* in Chicks. *Avian Diseases* 39:375-381.
55. Tschirdewahn, B., S. Notermans, K. Wernars, and F. Untermann. 1991. The Presence of Enterotoxigenic *Clostridium perfringens* Strains in Faeces of Various Animals. *International Journal of Food Microbiology* 14:175-178.
56. Wati, T., Ghosh, T., Syed, B., and S. Haldar. 2015. Comparative efficacy of a phytogenic feed additive and an antibiotic growth promoter on production performance, caecal microbial population and humoral immune response of broiler chickens inoculated with enteric pathogens. *Animal Nutrition* 1(2015): 213-219.
57. Williams, R.B. 2005. Intercurrent Coccidiosis and Necrotic Enteritis of Chickens: Rational, Integrated Disease Management by Maintenance of Gut Integrity. *Avian pathology* 34(3):159-180.
58. Williams, R. B., R. N. Marshall, R. M. La Regione, and J. Catchpole. 2003. A New Method for the Experimental Production of Necrotic Enteritis and its Use for Studies on the Relationships Between Necrotic Enteritis, Coccidiosis and Anticoccidial Vaccination of Chickens. *Parasitology Research* 90:19-26.
59. Wise, M. G., and G. R. Siragusa. 2005. Quantitative Detection of *Clostridium perfringens* in the Broiler Fowl Gastrointestinal Tract by Real-Time PCR. *Applied and Environmental Microbiology*

71:3911-3916.

60. Wiseman, R.W., Bushnell, O.A., and Rosengerg, M.M. 1956. Effects of Rations on the pH and Microflora in Selected Regions of the Intestinal Tract of Chickens. *Poultry Science* 35:126-132.
61. Yegani, M., and D. R. Korver. 2008. Factors Affecting Intestinal Health in Poultry. *Poultry Science* 87:2052-2063.