

Necrotic enteritis: The complete overview



by **Inge Heinzl** and **Technical Team**, EW Nutrition

Necrotic enteritis is a profit killer in poultry production

Necrotic enteritis is the cause of USD 6 billion losses every year in global poultry production, corresponding to USD 0.0625 per bird (Wade and Keyburn, 2015). This controllable disease is on the rise. One reason is the voluntary or legally required reduction of antibiotics in animal production due to the increasing occurrence of antimicrobial resistance but also consumer demand. Another reason is the administration of live Coccidiosis vaccines and partial reduction of ionophores, which also show efficacy against Gram-positive bacteria (Williams, 2005).

Necrotic enteritis and coccidiosis are the most significant health problem in broilers (Hofacre et al., 2018).

The disease generally occurs in broiler chickens of 2-6 weeks of age. It is caused by an overgrowth of *Clostridium perfringens* type A and, to a lesser extent, type C in the small intestine. The toxins produced by *C. perfringens* also damage the intestinal wall.

Clinical and subclinical forms of NE - which one causes more significant losses?

The clinical form is obvious



Intestine showing signs of NE

...is characterized by acute, dark diarrhea resulting in wet litter and suddenly increasing flock mortality of up to 1% per day after the first clinical signs appear (Ducatelle and Van Immerseel, 2010), sometimes summing up to mortality rates of 50% (Van der Sluis, 2013). The birds have ruffled feathers, lethargy, and inappetence.

Necropsy typically shows ballooned small intestines with a roughened appearing mucosal surface, lesions, and brownish (diphtheritic) pseudomembranes. There is a lot of watery brown, blood-tinged fluid and a foul odor during post-mortem examination. The liver is dark, swollen, and firm, and the gall bladder is distended (Hofacre et al., 2018).

In the case of **peracute** Necrotic Enteritis, birds may die without showing any preliminary signs.

The subclinical form often only can be noticed at the end of the cycle

When birds suffer from the subclinical form, chronic damage to the intestinal mucosa and an increased quantity of mucus in the small intestine lead to impaired digestion and absorption of nutrients resulting in poor growth performance. The deteriorated feed conversion and the resulting decreased performance become noticeable around day 35 of age. As feed contributes approximately 65-75% of the input cost to produce a broiler chicken, poor feed conversion increases production costs and significantly influences profitability. Often, due to a lack of clear symptoms, this subclinical disease remains untreated and permanently impacts the efficiency of production.



The pathogen causing NE - a ubiquitous bacterium

Responsible for Necrotic Enteritis are Gram-positive, anaerobic bacteria, specific strains of *Clostridium perfringens* type A and, to a lesser extent, type C (Keyburn et al., 2008).



Clostridia primarily occur in the soil where organic substances are degraded, in sewage, and in the gastrointestinal tract of animals and humans. These bacteria produce spores, which are extremely resistant to environmental impact (heat, irradiation, exsiccation), some disinfectants, and can survive for several years. Under suitable conditions, *C. perfringens* spores can even proliferate in feed or litter.

Clostridium perfringens is a “natural inhabitant” of the intestine of chickens. In healthy birds, it occurs in a mixture of diverse strains at 10^2 - 10^4 CFU/g of digesta (McDevitt et al., 2006). The disease starts when *C. perfringens* proliferates in the small intestine, usually due to a combination of factors such as high amount protein, low immunity, and an imbalance in the gut flora. Then, the number rises to 10^7 - 10^9 CFU/g of digesta (Dahiya et al., 2005).

Highly important: NetB, a pore-forming toxin is a key virulence factor for NE

To establish in the host, *Clostridium* Spp. and other pathogens depend on virulence factors (see infobox). These virulence factors include for example “tools” for attachment, evasion or suppression of the host’s immune system, “tools” for getting nutrients, and “tools” for entry into intestinal cells. Over the years, the α -toxin produced by *C. perfringens* was assumed to be involved in the development of the disease and a key virulence factor. In 2008, Keyburn and coworkers found another key virulence factor by using a *C. perfringens* mutant unable to produce α -toxin, while still causing Necrotic Enteritis.

Thus, another toxin was identified occurring only in chickens suffering from Necrotic Enteritis: *C. perfringens* necrotic enteritis B-like toxin (NetB). NetB is a pore-forming toxin. Pore-forming toxins are exotoxins usually produced by pathogenic bacteria but may also be produced by other microorganisms. These toxins destroy the integrity of gut wall cell membranes. The leaking cell contents serve as nutrients for the bacteria. If immune cells are destroyed, an immune reaction might be partially impaired.

Additionally, pathogenic strains of *C. perfringens* produce bacteriocins – the most important is Perfrin (Timbermont et al., 2014) – to inhibit the proliferation of harmless *Clostridium* Spp. strains and to replace the normal intestinal flora of chickens (Riaz et al., 2017).

Examples of virulence factors

1. Adhesins

Enable the pathogen to adhere or attach within the target host site, e.g. via fimbria. Pili enable the exchange of RNA or DNA between pathogens.

2. Invasion factors


Facilitate the penetration and the distribution of the pathogens in the host tissue (invasion and spreading enzymes). For example: hyaluronidase attacking the hyaluronic acid of the connective tissue or flagella enabling the pathogens to actively move.

3. Toxins

Damage the function of the host cells or destroy them (e.g. endotoxins – lipopolysaccharides, exotoxins)

4. Strategies of evasion

Enable the pathogen to undergo the strategies of defense of the immune system (e.g. antiphagocytosis factors provide protection against an attack by phagocytes; specific antibodies are inactivated by enzymes).



Predisposing factors favor the development of NE

A chicken with **optimal gut health** may be less **susceptible to NE**. Additional **predisposing factors** are necessary to allocate nutrients and make the gut environment suitable for the proliferation of these pathogens, enabling them to cause disease (Van Immerseel et al., 2008; Williams, 2005).

1. FEED: composition and particle size are critical

Feed plays a role in the development of Necrotic Enteritis that should not be underestimated. Here, substances creating an intestinal environment favorable for *C. perfringens* must be mentioned.

NSPs

Non-starch polysaccharides

Kaldhusdal and Skjerve (1996); Annett et al. (2002)

Mode of action:

- Increase viscosity
- Decrease passage rate
- Decrease digestibility of other nutrients
- Serve as nutrient



more nutrients available for
C. perfringens

- Stimulation of mucus production



anaerobic environment
Van der Sluis, 2013

Protein and fat

Excessive content of protein

Poorly digestible protein

especially protein (fish meal) with high
Concentrations of zinc/glycine



shift in balance of intestinal flora
(Antonissen et al., 2016)

Mode of action:

- High protein levels in the GIT
- Act as substrate for the bacteria


Animal fat

Mode of action:

- Higher pH and higher bile salt conc.
- Promotion of coccis



increased growth of *C. perfringens*
(Knarreborg et al., 2002)



2. Mycotoxins create ideal conditions for NE

Mycotoxins harm gut integrity and create ideal conditions for the proliferation of *Clostridium perfringens*:

Mycotoxins do not have a direct effect on *C. perfringens* proliferation, toxin production, or NetB transcription. However, mycotoxins disrupt gut health integrity, creating a favorable environment for the pathogen. For example:

- DON provides good conditions for proliferation of *perfringens* by disrupting the intestinal barrier and damaging the epithelium. The possibly resulting permeability of the epithelium and a decreased absorption of dietary proteins can lead to a higher amount of proteins in the small intestine. These proteins may serve as nutrients for the pathogen (Antonissen et al., 2014).
- DON and other mycotoxins decrease the number of lactic acid producing bacteria indicating a shift in the microbial balance (Antonissen et al., 2016.)

3. Eimeria spp.: forming a perfect team with Clostridium perfringens

An intact intestinal epithelium is the best defense against potential pathogens such as *C. perfringens*. Here, coccidiosis comes into play. Moore (2016) showed that by damaging the gut epithelium, *Eimeria* species give *C. perfringens* access to the intestinal basal domains of the mucosal epithelium. Then, the first phase of the pathological process takes place and from there, *C. perfringens* invades the lamina propria. Damage to the epithelium follows (Olkowski et al., 2008). The plasma proteins leaking to the gut and the mucus produced are rich nutrient sources (Van Immerseel et al., 2004; Collier et al., 2008). A further impact of coccidiosis is shifting the microbial balance in the gut by decreasing the number of e.g., *Candidatus savagella* which activates the innate immune defense.




Figure 1

Figure 1:

- A. *Eimeria* induce leakage of plasma proteins by killing epithelial cells
- B. They enhance mucus production in the intestine

A+B lead to an increase in available nutrients and create an environment favorable for the proliferation of *perfringens*

Not only *Eimeria* Spp., also other pathogens (e.g. *Salmonella* Spp., Ascarid larvae, viruses) and agents, such as mycotoxins damaging the intestinal mucosa can pave the way for a *C. perfringens* infection. Predisposing factors like wet litter, the moisture of which is essential for the sporulation of *Eimeria* Spp. oocysts, must also be considered as promoting factors for Necrotic Enteritis (Williams, 2005).

4. Immunosuppressive Factors: Bacteria, viruses..., and stress

Any factor which induces stress in the animals disrupts the balance of the intestinal flora. The resulting suppression of the immune system contributes to the risk of Necrotic Enteritis (Tsiouris, 2016).

Bacteria

Shivaramaiah and coworkers (2011) investigated a neonatal *Salmonella typhimurium* infection as a predisposing factor for NE. The early infection causes significant damage to the gut (Porter et al., 1998). Additionally, Hassan et al. (1994) showed that the challenge with *Salmonella typhimurium* negatively impacted the development of lymphocytes which might also promote a colonization of *Clostridium perfringens*.

Viruses

Infectious Bursal Disease is known to increase the severity of infections with salmonella, staphylococci, but also clostridia. Another clostridia-promoting viral disease is Marek's Disease.

Stress:

The intestinal tract is particularly sensitive to any type of stress. This stress can be caused by e.g. too high temperatures, high stocking densities, an abrupt change of feed.




Figure 2: Predisposing factors weakening the birds and enabling *Clostridium* to attack

Treatment is necessary in the case of acute disease

In this instance, the farmer is obligated to consult a veterinarian and treat his birds.

It must be mentioned that, as the treatment takes place via feed or water, only birds which still consume water or feed may be treated.

Antibiotics are effective but also take a risk

Antibiotics targeting Gram-positive bacteria are commonly used for the treatment of acute NE. The antibiotic choice shall be addressed by a veterinarian, taking into account the mode of action and the presence of resistance genes in the farm/flock.

The prophylactic use of antibiotics is not recommended and many countries have already [banned it in order to reduce antimicrobial resistance \(AMR\)](#).

Antimicrobial Resistance (AMR)

Some bacteria are less sensitive to certain antibiotics due to genetic mutations. They are able to:

- stimulate the production of enzymes, which break down or modify the antibiotics and inactivate them (1).
- eliminate entrances for antibiotics or promote the development of pumps, which discharge the antibiotic before taking effect (2).
- change or eliminate molecules to which the antibiotic would bind (targets for the antibiotics).

This means that, when the corresponding antibiotics are used, bacteria resistant against these antibiotics survive. Due to the fact that their competitors have been eliminated they are able to reproduce better.

Additionally, this resistance may be transferred by means of “resistance genes”

- to daughter cells
- via their intake from dead bacteria (3)
- through horizontal gene transfer (4)
- through viruses (5)

Every application of antibiotics promotes the development of resistance (Robert Koch Institute, 2019). A short-term use, [better biosecurity](#), or an application at low dosage give the bacteria a better chance to adapt.

Bacteriophages would be possible but are still disputed

Experimental use of phage treatments has shown to be effective in reducing disease progression and symptoms of Necrotic Enteritis (Miller et al., 2010). By oral application of a bacteriophage cocktail, Miller and coworkers could reduce mortality by 92% in *C. perfringens*-challenged broilers compared to the untreated control.

Mode of action: the endolysins, highly evolved enzymes produced by bacteriophages, are able to digest the bacterial cell wall for phage progeny release (Fischetti, 2010). However, phages are still not approved by the EFSA.

Excurs:

Antimicrobial Resistance (AMR) Some bacteria are less sensitive to certain antibiotics due to genetic mutations. They are able to:

- stimulate the production of enzymes, which break down or modify the antibiotics and inactivate them (1).
- eliminate entrances for antibiotics or promote the development of pumps, which discharge the antibiotic before taking effect (2).
- change or eliminate molecules to which the antibiotic would bind (targets for the antibiotics) (figure 3)




Figure 3: Possibilities of a bacterial cell to defend itself against antibiotics

This means that, when the corresponding antibiotics are used, bacteria resistant against these antibiotics survive. Due to the fact that their competitors have been eliminated they are able to reproduce better. Additionally, this resistance may be transferred by means of “resistance genes”

- to daughter cells
- via their intake from dead bacteria (3)
- through horizontal gene transfer (4)
- through viruses (5) (figure 4)




Figure 4: Possibilities to transfer resistance to other bacterial cells

Every application of antibiotics promotes the development of resistance (Robert Koch Institute, 2019). A short-term use or an application at low dosage give the bacteria a better chance to adapt.


Preventing a disease is always better than its treatment!

But how to do it? Preventing the conditions that favor the proliferation of *Clostridium perfringens* and strengthening the host's immune response lowers the probability of disease. Besides eliminating the predisposing factors, the main targets are:

- **Balance of the gut flora**
- **Optimization of gut function and integrity**
- **Maintenance of immunity**

1. Biosecurity is of the highest importance!

There is evidence that most *Clostridium* strains isolated from birds suffering from Necrotic Enteritis could induce the disease experimentally, while strains isolated from healthy birds cannot. This confirms that only specific strains are problematic (Ducatelle and Van Immerseel, 2010). So, it's of highest importance to avoid introducing these pathogenic strains to the farm.



Strict biosecurity measures!

Separate clothing, boots, and hand washing/disinfecting facilities in each poultry house

More than 14 days of down time between flocks

2. Specific measures against coccidiosis

1. Vaccination

According to parasitologists, 7 to 9 *Eimeria* species are found in chickens, and they do not cross-protect against each other. An effective vaccination must contain sporulated oocysts of the most critical pathogenic *Eimeria* species (*E. acervulina*, *E. maxima*, *E. tenella*, *E. necatrix*, and *E. brunetti*). The more species contained in the vaccine, the better. However, if not applied the correct way, vaccines can be ineffective or cause reactions in the birds that might lead to NE (Mitchell, 2017).

2. Anticoccidials

Alternate use of chemicals (synthetic compounds) and ionophores (polyether antibiotics) with different modes of action is important to avoid development of resistance.

Ionophores have a specific mode of action and kill oocysts before they are able to infect birds. Being very small, ionophore molecules can be taken up and diffused into the outer membrane of the sporozoite. There, it decreases the concentration gradient leading to an accumulation of water within the sporozoite causing its bursting.

3. Diet – favorable for the birds but not for clostridium!

Minimizing non-starch polysaccharides (NSPs) in cereals



To prevent a “feeding” of *Clostridium perfringens*, high content of water-soluble but indigestible NSPs such as wheat, wheat by-products, and barley should be avoided or at least minimized. Additionally, xylanases should be included in the feed formulation to reduce the deleterious effects of NSPs and improve feed energy utilization. Instead of these cereals, maize could be included in the diet. It is considered a perfect ingredient in broiler diets due to its high energy content and high nutrient availability.

Formulating low protein diets/diets with highly digestible amino acids

Feeding low-protein diets supplemented with crystalline amino acids might be beneficial to reduce the risk of Necrotic Enteritis (Dahiya et al., 2007). To improve protein digestibility and therefore reduce the proliferation of *C. perfringens*, proteases may be added to the feed.

Avoiding/Minimizing animal fats in the diet

Animal fats tend to increase the counts of *Clostridium perfringens*; thus, they should be replaced by vegetable fat sources.

Feed form is decisive

In terms of feed form, Engberg et al. (2002) found that birds fed pellets showed a reduced number of *Clostridium perfringens* in the caeca and the rectum than mash-fed birds. Branton and coworkers (1987) reported a lower mortality by feeding roller-milled (coarsely ground) than hammer-milled feed.

4. Additives

Additives can be used either to prevent the proliferation of *Clostridium perfringens* or to change the environmental conditions in a way that proliferation of *C. perfringens* is prevented.

1. Probiotics directly support the balance of the microbiome

These live microbial supplements can be used to help to establish, maintain or re-establish the intestinal microflora.

Mode of action:

They compete with pathogenic bacteria for substrates and attachment sites and produce antimicrobial substances inhibiting the growth of pathogenic bacteria (Gillor et al., 2008). They bind and neutralize enterotoxins (Mathipa and Thantsha, 2017) and promote immune function of the host (Yang et al., 2012)

2. Prebiotics indirectly promote the microbiome

These feed ingredients serve as substrates to promote beneficial bacteria in the intestine.

Mode of action:

D-mannose or fructose, starches non-digestible by birds, selectively stimulate the growth and the activity of the “good” gut flora. Fructooligosaccharides decrease *C. perfringens* and *E. coli* in the gut and increase the diversity of *Lactobacillus* Spp. (Kim et al., 2011). Galactooligosaccharides, in combination with a *B. lactis*-based probiotic, have been reported to selectively promote the proliferation of *Bifidobacterium* Spp. (Jung et al., 2008).

3. Organic acids support gut health

[Organic acids](#) are often used in animal diets to improve intestinal health.

Mode of action:

A decreased pH promotes beneficial bacteria. Caprylic acid suppresses *C. perfringens* but also *Salmonella* spp. by inhibiting their utilization of glucose (Skrivanova et al., 2006). Lauric, citric, oleic, and linoleic acid, as well as medium-chain fatty acids (C8-C14), impede the growth of *C. perfringens*.

A trial with different organic acid products showed high efficacy for Acidomix AFG and Acidomix AFL against *Clostridium perfringens* as well as against *Salmonella enterica*. For the test, 50 µl solution containing different microorganisms (reference strains of *S. enterica* and *C. perfringens*; conc. 10⁵ CFU/ml) together with 50 µl of increasing concentrations of various organic acids/organic acid products (Acidomix) were pipetted into microdilution plates. After the respective incubation, the MICs of every organic acid/organic acid product were calculated.

Figure 5 shows the minimum inhibiting concentrations (MIC). For Acidomix AFL and AFG, lower concentrations than for fumaric, lactic, and propionic acid were needed to inhibit the growth of *Salmonella enterica* and *Clostridium perfringens*.




Figure 5: Minimal inhibiting concentrations of Acidomix AFL and Acidomix AFG against *Salmonella enterica* and *Clostridium perfringens*

Phytomolecules : different types are available against NE

[Phytomolecules](#), also known as secondary plant compounds, have been used against pathogens for centuries. In general, two subgroups of these substances are known as effective against *Clostridium perfringens*: Tannins and Essential Oils.

Tannins

Many studies have shown the efficacy of [tannins](#) against different pathogens such as helminths, *Eimeria* spp., viruses, and bacteria. Extracts from the chestnut and quebracho trees are effective not only against *C. perfringens* but also its toxins (Elizando et al., 2010). Tannins act against *Eimeria* spp. (Cejas et al., 2011) and *Salmonella* Sp., two predisposing factors for NE.

A trial was conducted with Pretect D, a product based on tannins and saponins, to show its efficacy against coccidia, one of the predisposing factors of NE. For the 35-day study conducted at a commercial research facility in the US, 1800 one-day-old Cobb 500 broilers were divided into four groups of 450 birds each (with 9 replicates & 50 birds per replicate). They all received the standard feed of the farm (Starter D0-D21, Grower D22-D35).

The challenge was given in the form of a freshly prepared mixed inoculum with *E. acervulina* (100,000 oocysts/ bird), *E. maxima* (50,000 oocysts/ bird), and *E. tenella* (75,000 oocysts/ bird). The inoculum was mixed into the feed in the base of each pen's tube feeder.

The oocyst count per gram of feces (OPG) was done on D21, D27 & D35. The cocci lesion scoring (CLS) was done on D27 following Johnson and Reid (1970) with 0=normal; 4=most

Group	Challenge	Additive
Non-challenged Control (NC)	No	No
Challenged Control (CC)	Yes	No
CC + Ionophore	Yes	Ionophore@60ppm
CC + Pretect D	Yes	Pretect D@500ppm

The trial showed that, due to Pretect D, the lesion score showed a lower value indicating that lesions could be reduced or were less severe, which can be seen in figure 6:




Figure 6: Average lesion score

Essential Oils

Their hydrophobic characteristic enables them to interact with the lipids of the membrane of *C. perfringens*. They can incorporate into the bacterial membrane and disrupt its integrity, increasing the permeability of the cell membrane for ions and other small molecules such as ATP and leading to the decrease of the electrochemical gradient above the cell membrane and the loss of the cell's energy equivalents. Besides their direct effect on *Clostridium spp.*, many phytomolecules improve gut health and help prevent the proliferation of *Clostridium spp.* And, therefore, Necrotic Enteritis.

An In vitro-trial shows Ventar D reducing Clostridia and sparing the beneficial lactobacilli. In this trial, the bacteria (*Clostridium perfringens*, *Lactobacillus agilis* S73, and *Lactobacillus plantarum*) were cultured under favorable conditions (RCM, 37°C, anaerobe for *Clostridium perfringens*, and MRS, 37°C, 5 % CO₂ for Lactobacilli) and exposed to different concentrations of Ventar D (0 µg/ml – control, 500 µg/ml, 750 µg/ml, and 1000 µg/ml).

The results of the trial with *Clostridium perfringens* are shown in figure 7.




Figure 7: Different concentrations of Ventar D added to *Clostridium perfringens* cultures

Here, a significant reduction of colonies could already be observed at a concentration of 500 µg/ml of Ventar D. With 750 µg/ml, only a few colonies remained, and at a Ventar D concentration of 1000 µg/ml, *Clostridium perfringens* didn't grow anymore.

In contrast, the Lactobacilli showed a different picture: only at the higher concentration (1250 µg/ml of Ventar D) did *Lactobacillus plantarum* and *Lactobacillus agilis* S73 show a slight growth reduction (figure 8).





Figure 8: *Lactobacillus plantarum* exposed to 0 (left) and 1250 µg/ml (right) of Ventar D

1. In vivo-trial in poultry shows that phytomolecules reduce gut lesions

The study was conducted at Southern Poultry Feed & Research, Athens, GA (USA), over 42 days. It included in total 880 Cobb 500 broilers across 2 trial groups, with 11 repetitions per trial set-up and 40 animals per replicate floor pen. All animals received routine vaccinations at the hatchery and were healthy when starting the trial.

Control group	Built-up litter (no additive)
Ventar D group	Built-up litter + Ventar D, 100 g/MT of feed


All birds received standard feed, fed as crumbles/pellets ad libitum. Feed intake by pen was recorded per feeding phase for starter (D21), grower (D35), and finisher feed (D42). Bird weights were recorded at study initiation, on D21, D35, and D42. On D21 and D35, three birds per pen were sacrificed. The GIT was scored for necrotic enteritis lesions; figures 9 and 10 show the results.



Figures 9 and 10: Lesion score on days 21 and 35

Already on day 21, the birds of the Ventar D group showed a less impacted gut mucosa, indicated by a lower lesion score. Lesions were reduced in both groups until day 35; however, the value of the Ventar D group was still better.

A less impacted gut has a higher digestion and absorption capacity, which results in better performance (FCR and weight gain) and lower mortality (figures 11-14).



Figures 11-14: Performance data of a control group compared with birds supplemented with Ventar D

The two trials show that Ventar D allows the poultry producer to proactively strengthen broilers' gut health by controlling *Clostridia perfringens* and promoting/saving beneficial bacteria such as lactobacilli. The effects of the reduction of *Clostridia* can be seen in vivo in a lower lesion score and better performance.

Toxin binders adsorb bacterial and mycotoxins

These binders have two modes of action:

They bind mycotoxins and, therefore, reduce or prevent damage to the intestinal wall so that the preconditions for *Clostridium* spp. proliferation are not generated.

Additionally, binding toxins produced by *Clostridium perfringens* can reduce the occurrence or severity of lesions: Alpha-toxin, phospholipase C, hydrolyses membrane phospholipids and damages erythrocytes, leucocytes, myocytes, and endothelial cells and causes their lipolysis (Songer, 1996), leading to necrosis and tissue damage.

Binding NetB toxin, the key virulence factor, could reduce the severity of Necrotic Enteritis.

A trial was conducted in a laboratory in Valladolid/Spain to show the high binding capacity of Solis Plus 2.0. All tests were carried out as duplicates and using a standard liquid chromatography/mass spectrometry (LC/MS/MS) quantification. Interpretation and data analysis were carried out with the corresponding software. Toxin concentrations, anti-mycotoxin agent application rates, and pH levels were set as follows:

Mycotoxin	Challenge Level	Challenge (ppb)	Solis Plus 2.0 inclusion rate	Assay time
Aflatoxin	Low	150	0.2%	30 min.
	High	1500		
Fumonisin	Low	500		
	High	5000		
Ochratoxin	Low	150		
	High	1500		

The results are shown in figure 15:




Figure 15: Adsorption capacity of Solis Plus for relevant mycotoxins

Under acidic conditions (pH 3), Solis Plus 2.0 effectively adsorbs the three tested mycotoxins at low and high contamination levels:

- Aflatoxin: 150 ppb -100 %; 1500 ppb – 98 %
- Fumonisin: 500 ppb – 87%; 5000 ppb – 86 %
- Ochratoxin: 150 ppb – more than 43 %; 1500 ppb – 52 %.

By binding harmful toxins and preventing their negative impact on the gut, toxin binders can also be a tool to reduce necrotic enteritis.

NE can be controlled - even in an antibiotic-free era

The ever-growing trend of reduced antibiotic and ionophore use increases the incidence of Necrotic Enteritis in poultry production. Especially the subclinical form, which generally goes unnoticed, results in poor feed efficiency and is a major cause of financial losses to poultry producers.

Maintaining optimum gut health is key to preventing the occurrence of Necrotic Enteritis. In the era of antibiotic-free poultry production, alternatives acting against the pathogenic bacterium and also against its predisposing factors must be considered to control this devastating disease. The industry already provides solutions like phytomolecules-based products or toxin binders to support the animals.

References:

Annett, C.B., J. R. Viste, M. Chirino-Trejo, H. L. Classen, D. M. Middleton, and E. Simko. "Necrotic enteritis: effect of barley, wheat and corn diets on proliferation of *Clostridium perfringens* type A." *Avian Pathology* 31 (2002): 599– 602. <https://doi.org/10.1080/0307945021000024544>

Antonissen G, F. Van Immerseel, F. Pasmans, R. Ducatelle, F. Haesebrouck, L. Timbermont, M. Verlinden, G.P.J.

- Janssens, V. Eeckhaut, M. Eeckhout, S. De Saeger, S. Hessenberger, A. Martel, and S. Croubels. "The mycotoxin deoxynivalenol predisposes for the development of *Clostridium perfringens*-Induced necrotic enteritis in broiler chickens. *PLoS ONE* 9 no. 9 (2014): e108775. <https://doi.org/10.1371/journal.pone.0108775>
- Antonissen, G., V. Eeckhaut, K. Van Driessche, L. Onrust, F. Haesebrouck, R. Ducatelle, R.J. Moore, and F. Van Immerseel. "Microbial Shifts Associated With Necrotic Enteritis." *Avian Pathol.* 45 no. 3 (2016): 308-312. <https://doi.org/10.1080/03079457.2016.1152625>
- Branton, S.L., F.N. Reece, and W.M. Hagler. "Influence of a wheat diet on mortality of broiler chickens associated with necrotic enteritis." *Poultry Sci.* 66 (1987): 1326-1330. <https://doi.org/10.3382/ps.0661326>
- Cejas, E., S. Pinto, F. Prosdócimo, M. Batalle, H. Barrios, G. Tellez, and M. De Franceschi. "Evaluation of quebracho red wood (*Schinopsis lorentzii*) polyphenols vegetable extract for the reduction of coccidiosis in broiler chicks." *International Journal of Poultry Science* 10 no. 5 (2011): 344-349. <https://doi.org/10.3923/ijps.2011.344.349>
- Collier, C.T., C.L. Hofacre, A.M. Payne, D.B. Anderson, P. Kaiser, R.I. Mackie, and H.R. Gaskins. "Coccidia-induced mucogenesis promotes the onset of necrotic enteritis by supporting *Clostridium perfringens* growth." *Veterinary Immunology and Immunopathology* 122 (2008):104-115. <https://doi.org/10.1016/j.vetimm.2007.10.014>
https://www.academia.edu/12692646/Coccidia-induced_mucogenesis_promotes_the_onset_of_necrotic_enteritis_by_supporting_Clostridium_perfringens_growth
- Dahiya, J.P., D. Hoehler, A.G. Van Kessel, and M.D. Drew. "Effect of different dietary methionine sources on intestinal microbial populations in broiler chickens." *Poultry Science* 86 (2007):2358-2366 <https://doi.org/10.3382/ps.2007-00133>
- Dahiya, J.P., D. Hoehler, D.C. Wilkie, A.G. van Kessel, and M.D. Drew. "Dietary glycine concentration affects intestinal *Clostridium perfringens* and *Lactobacilli* populations in broiler chickens." *Poultry Science* 84 no.12 (2005):1875-85. <https://doi.org/10.1093/ps/84.12.1875>
- Diaz Carrasco, J.M., L.M. Redondo, E.A. Redondo, J.E. Dominguez, A.P. Chacana, and M.E. Fernandez Miyakawa. "Use of plant extracts as an effective manner to control *Clostridium perfringens* induced necrotic enteritis in poultry." *BioMed Research International* (2016): Article ID 3278359. <https://dx.doi.org/10.1155/2016/3278359>
- Ducatelle, R. and F. van Immerseel. "Necrotic enteritis: emerging problem in broilers." *WATTAgNet.com - Poultry Health and Disease* (April 9, 2010). <https://www.wattagnet.com/articles/5523-necrotic-enteritis-emerging-problem-in-broilers>
- Elizondo, A.M., E.C. Mercado, B.C. Rabinovitz, and M.E. Fernandez-Miyakawa. "Effect of tannins on the in vitro growth of *Clostridium perfringens*." *Veterinary Microbiology* 145 no. 3-4 (2010): 308-314. <https://doi.org/10.1016/j.vetmic.2010.04.003>
- Engberg, R.M., M.S. Hedemann, and B.B. Jensen. "The influence of grinding and pelleting of feed on the microbial composition and activity in the digestive tract of broiler chickens." *British Poultry Science* 43 no. 4 (2002):569-579. <https://doi.org/10.1080/0007166022000004480>
- Fischetti, V.A. "Bacteriophage endolysins: A novel anti-infective to control Gram-positive pathogens." *J Med Microbiol.* 300 no. 6 (2010): 357-362. <https://doi.org/10.1016/j.jimm.2010.04.002>
- Gillor, O., A. Etzion and M.A. Riley. "The dual role of bacteriocins as anti- and probiotics." *Appl Microbiol Biotechnol.* 81 no. 4 (2008): 591-606. <https://doi.org/10.1007/s00253-008-1726-5>
- Hofacre, C.L., J.A. Smith, and G.F. Mathis. "Invited Review. An optimist's view on limiting necrotic enteritis and maintaining broiler gut health and performance in today's marketing, food safety, and regulatory climate." *Poultry Science* 97 (2018):1929-1933. <https://dx.doi.org/10.3382/ps/pey082>
- Jung, S.J., R. Houde, B. Baurhoo, X. Zhao, and B. H. Lee. "Effects of galacto-oligosaccharides and a bifidobacteria lactis-based probiotic strain on the growth performance and fecal microflora of broiler chickens." *Poultry Science* 87 (2008):1694-1699. <https://doi.org/10.3382/ps.2007-00489>

Kaldhusdal and Skjerve. "Association between cereal contents in the diet and incidence of necrotic enteritis in broiler chickens in Norway." *Preventive Veterinary Medicine* 28 (1996):1-16.

[https://doi.org/10.1016/0167-5877\(96\)01021-5](https://doi.org/10.1016/0167-5877(96)01021-5)

https://www.academia.edu/17521917/Association_between_cereal_contents_in_the_diet_and_incidence_of_necrotic_enteritis_in_broiler_chickens_in_Norway

Keyburn, A. L., S. A. Sheedy, M. E. Ford, M. M. Williamson, M. M. Awad, J. I. Rood, and R. J. Moore. "Alpha-toxin of *Clostridium perfringens* is not an essential virulence factor in necrotic enteritis in chickens." *Infect. Immun.* 74 (2006): 6496-6500. <https://doi.org/10.1128/IAI.00806-06>

Keyburn, A.L., J.D. Boyce, P. Vaz, T.L. Bannam, M.E. Ford, D. Parker, A. Di Rubbo, J.I. Rood, and R.J. Moore. "NetB, a new toxin that is associated with avian necrotic enteritis caused by *Clostridium perfringens*." *PLoS Pathog* 4 no. 2, e26 (2008): 0001-0011. <https://doi.org/10.1371/journal.ppat.0040026>

Kim, G.-B., Y. M. Seo , C. H. Kim , and I. K. Paik. "Effect of dietary prebiotic supplementation on the performance, intestinal microflora, and immune response of broilers." *Poultry Science* 90 (2011):75-82.

<https://doi.org/10.3382/ps.2010-00732>

Knap, I., B. Lund, A. B. Kehlet, C. Hofacre, and G. Mathis. "Bacillus licheniformis prevents necrotic enteritis in broiler chickens." *Avian Diseases* 54 no. 2 (2010):931-935. <https://doi.org/10.1637/9106-101509-ResNote.1>

Knarreborg, A., M.A. Simon, R.M. Engberg, B.B. Jensen, and G.W. Tannock. "Effects of Dietary Fat Source and Subtherapeutic Levels of Antibiotic on the Bacterial Community in the Ileum of Broiler Chickens at Various Ages." *Applied and Environmental Microbiology* 68 no. 12 (2002): 5918-5924.

<https://doi.org/10.1128/AEM.68.12.5918-5924.2002>

Kocher, A. and M. Choct. "Improving broiler chicken performance. The efficacy of organic acids, prebiotics and enzymes in controlling necrotic enteritis." *Australian Government-Rural Industries Research and Development Corporation*. Publ. no. 08/149 (2008).

<https://www.agrifutures.com.au/wp-content/uploads/publications/08-149.pdf>

Kondo, F. "In vitro lecithinase activity and sensitivity to 22 antimicrobial agents of *Clostridium perfringens* isolated from necrotic enteritis of broiler chickens." *Research in veterinary Science* 45 (1988): 337-340.

[https://doi.org/10.1016/S0034-5288\(18\)30961-5](https://doi.org/10.1016/S0034-5288(18)30961-5)

Kubena, L.F., J.A. Byrd, C.R. Young, and D.E. Corrier. "Effects of tannic acid on cecal volatile fatty acids and susceptibility to *Salmonella typhimurium* colonization in broiler chicks." *Poultry Science* 80, no. 9, pp. 1293-1298, 2001.

<https://doi.org/10.1093/ps/80.9.1293>

M'Sadeq S.A., Shubiao Wu, Robert A. Swick, Mingan Choct. "Towards the control of necrotic enteritis in broiler chickens with in-feed antibiotics phasing-out worldwide." *Animal Nutrition* 1 (2015): 1-11.

<https://dx.doi.org/10.1016/j.aninu.2015.02.004>

Mathipa, M.G. and M.S. Thantsha. "Probiotic engineering: towards development of robust probiotic strains with enhanced functional properties and for targeted control of enteric pathogens." *Gut Pathog.* 9 no. 28 (2017).

<https://doi.org/10.1186/s13099-017-0178-9>

McDevitt, R.M., J.D. Brooker, T. Acamovic, and N.H.C. Sparks. "Necrotic enteritis, a continuing challenge for the poultry industry." *World's Poultry Science Journal* 62; World's Poultry Science Association (June 2006).

<https://doi.org/10.1079/WPS200593>

Miller, R.W., J. Skinner, A. Sulakvelidze, G.F. Mathis, and C.L. Hofacre. "Bacteriophage therapy for control of Necrotic Enteritis of broiler chickens experimentally infected with *Clostridium perfringens*." *Avian Diseases* 54 no. 1 (2010): 33-40. <https://doi.org/10.1637/8953-060509-Reg.1>

Mitsch, P., K. Zitterl-Eglseer, B. Köhler, C. Gabler, R. Losa, and I. Zimpernik. "The Effect of Two Different Blends of Essential Oil Components on the Proliferation of *Clostridium perfringens* in the Intestines of Broiler Chickens." *Poultry Science* 83 (2004):669-675. <https://doi.org/10.1093/ps/83.4.669>

Mitchell, A. "Choosing the right coccidiosis vaccine for layer and breeder chickens." *The Poultry Site* March 21 (2017). <https://thepoultrysite.com/articles/choosing-the-right-coccidiosis-vaccine-for-layer-and-breeder-chickens>

- Olkowski, A.A., C. Wojnarowicz, M. Chirino-Trejo, B. Laarveld, and G. Sawicki. "Sub-clinical necrotic enteritis in broiler chickens: Novel etiological consideration based on ultra-structural and molecular changes in the intestinal tissue." *Veterinary Science* 85 (2008): 543-553. <https://doi.org/10.1016/j.rvsc.2008.02.007>
- Pan, D. and Z. Yu. "Intestinal microbiome of poultry and its interaction with host and diet." *Gut Microbes* 5 no. 1 (2014): 108-119. <https://dx.doi.org/10.4161/gmic.26945>
- Robert Koch Institut. "Grundwissen Antibiotikaresistenz". https://www.rki.de/DE/Content/Infekt/Antibiotikaresistenz/Grundwissen/Grundwissen_inhalt.html#:~:text=Wenn%20ein%20neues%20Antibiotikum%20auf,%C3%BCberleben%20und%20vermehren%20sich%20weiter.
- Rougière, N. and B. Carré. "Comparison of gastrointestinal transit times between chickens from D + and D- genetic lines selected for divergent digestion efficiency." *Animal* 4 no. 11 (2010): 1861-1872. <https://doi.org/10.1017/S1751731110001266>
- Santos, F.B.O., B.W. Sheldon, A.A. Santos Jr., and P.R. Ferket. "Influence of housing system, grain type, and particle size on Salmonella colonization and shedding of broilers fed triticales or corn-soybean meal diets." *Poultry Science* 87 (2008): 405-420. <https://dx.doi.org/10.3382/ps.2006-00417>
- Schiavone, A. , K. Guo, S. Tassone, L. Gasco, E. Hernandez, R. Denti, and I. Zoccarato. "Effects of a Natural Extract of Chestnut Wood on Digestibility, Performance Traits, and Nitrogen Balance of Broiler Chicks." *Poult Sci.* 87 no. 3 (2008): 521-527. <https://doi.org/10.3382/ps.2007-00113>
- Singh, Y., V. Ravindran, T.J. Wester, A.L. Molan, and G. Ravindran. "Influence of feeding coarse corn on performance, nutrient utilization, digestive tract measurements, carcass characteristics, and cecal microflora counts of broilers." *Poultry Science* 93 (2014): 607-616. <https://dx.doi.org/10.3382/ps.2013-03542>
- Skrivanova, E., M. Marounek, V. Benda, and P. Brezina. "Susceptibility of Escherichia coli, Salmonella sp. and Clostridium perfringens to organic acids and monolaurin." *Veterinarni Medicina* 51 no. 3 (2006): 81-88. <https://doi.org/10.17221/5524-VETMED>
- Songer, J.G. "Clostridial Enteric Diseases of Domestic Animals." *Clinical Microbiology Reviews* 9 no. 2 (1996): 216-234. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC172891/pdf/090216.pdf>
- Stanley D., Wu S.-B., Rodgers N., Swick R.A., and Moore R.J. "Differential Responses of Cecal Microbiota to Fishmeal, Eimeria and Clostridium perfringens in a Necrotic Enteritis Challenge Model in Chickens." *PLoS ONE* 9 no. 8 (2014): e104739. <https://doi.org/10.1371/journal.pone.0104739>
- Tan, L., D. Rong, Y. Yang, and B. Zhang. "Effect of Oxidized Soybean Oils on Oxidative Status and Intestinal Barrier Function in Broiler Chickens." *Brazilian Journal of Poultry Science* 18 no. 2 (2018): 333-342. <https://dx.doi.org/10.1590/1806-9061-2017-0610>
- Tan, L., D. Rong, Y. Yang, and B. Zhang. "The Effect of Oxidized Fish Oils on Growth Performance, Oxidative Status, and Intestinal Barrier Function in Broiler Chickens." *J. Appl. Poult. Res.* 28 (2019): 31-41. <http://dx.doi.org/10.3382/japr/pfy013>
- ThePoultrySite. "Necrotic Enteritis. Disease Guide". <https://thepoultrysite.com/disease-guide/necrotic-enteritis>
- Timbermont L., A. Lanckriet, J. Dewulf, N. Nollet, K. Schwarzer, F. Haesebrouck, R. Ducatelle, and F. Van Immerseel. "Control of Clostridium perfringens-induced necrotic enteritis in broilers by target-released butyric acid, fatty acids and essential oils." *Avian Pathol.* 39 no. 2 (2010): 117-21. <https://doi.org/10.1080/03079451003610586>
- Tsiouris, V. "Poultry management: a useful tool for the control of necrotic enteritis in poultry." *Avian Pathol.* 45 no. 3 (2016):323-325. <https://doi.org/10.1080/03079457.2016.1154502>
- Van der Most, P.J., B. de Jong, H.K. Parmentier and S. Verhulst. "Trade-off between growth and immune function: a meta-analysis of selection experiments." *Functional Ecology* 25 (2011): 74-80. <https://doi.org/10.1111/j.1365-2435.2010.01800.x>
- Van der Sluis, W. "Clostridial enteritis is an often underestimated problem." *Worlds Poult. Sci. J.* 16 (2000):42-43.
- Van der Suis, W. "Necrotic enteritis kills birds and profits." *Poultry World* Apr5 (2013).

<https://www.poultryworld.net/Health/Articles/2013/4/Necrotic-enteritis-kills-birds-and-profits-1220877W/>

Van Immerseel, F., J. De Buck, F. Pasmans, G. Huyghebaert, F. Haesebrouck, and R. Ducatelle. "Clostridium perfringens in poultry: an emerging threat of animal and public health." *Avian Pathology* 33 (2004): 537-549. <https://doi.org/10.1080/03079450400013162>

Van Immerseel, F., J.I. Rood, R.J. Moore, and R.W. Titball. "Rethinking our understanding of the pathogenesis of necrotic enteritis in chickens." *Trends in Microbiology* 17 no. 1 (2008):32-36. <https://doi.org/10.1016/j.tim.2008.09.005>


Wade, B., A. Keyburn. "The true cost of necrotic enteritis." *World Poultry* 31 no. 7 (2015): 16-17. <https://www.poultryworld.net/Meat/Articles/2015/10/The-true-cost-of-necrotic-enteritis-2699819W/>

Wade, B., A.L. Keyburn, T. Seemann, J.I. Rood, and R.J. Moore. "Binding of Clostridium perfringens to collagen correlates with the ability to cause necrotic enteritis in chickens." *Veterinary Microbiology* 180 no. 3-4 (2015): 299-303. <https://doi.org/10.1016/j.vetmic.2015.09.019>

Williams, R.B. "Intercurrent coccidiosis and necrotic enteritis of chickens: rational, integrated disease management by maintenance of gut integrity." *Avian Pathology* 34 no. 3 (2005):159-180. <https://doi.org/10.1080/03079450500112195>

Yang , C.M., G.T. Cao, P.R. Ferket, T.T. Liu, L. Zhou, L. Zhang, Y.P. Xiao, and A. G. Chen. "Effects of probiotic, Clostridium butyricum, on growth performance, immune function, and cecal microflora in broiler chickens." *Poultry Science* 91 (2012): 2121-2129. <https://dx.doi.org/10.3382/ps.2011-02131>

Antibiotic reduction: The increased importance of high-level biosecurity



Biosecurity is the foundation for all disease prevention programs (Dewulf, et al., 2018), and one of the most important points in antibiotic reduction scenarios. It includes the combination of all measures taken to reduce the risk of introduction and spread of diseases. It is based on the prevention of and protection against infectious agents by understating the disease transmission processes.

The application of consistently high standards of biosecurity can substantially contribute to the [reduction of antimicrobial resistance](#), not only by preventing the introduction of resistance genes into the farm but also by lowering the need to use antimicrobials (Davies & Wales, 2019).

Lower use of antimicrobials with higher biosecurity

Several studies and assessments relate that high farm biosecurity status and/or improvements in biosecurity lead to [reduced antimicrobial use](#) (Laanen, et al., 2013, Gelaude, et al., 2014, Postma, et al., 2016, Collineau, et al., 2017 and Collineau, et al., 2017a). Laanen, Postma, and Collineau studied the profile of swine farmers in different European countries, finding a relation between the high level of internal biosecurity, efficient control of infectious diseases, and reduced need for antimicrobials.

Reports on reduction on antibiotic use due to farm interventions are also available. Gelaude, et al. (2014), evaluated data from several Belgian broiler farms, finding a reduction of antimicrobial use by almost 30% within a year when biosecurity and other farm issues were improved. Collineau et al. (2017) studied pig farms in Belgium, France, Germany, and Sweden, in which the use of antibiotics was reduced on average by 47% across all farms. The researches observed that farms with the most strict biosecurity protocols, higher compliance, and who also took a multidisciplinary approach (making other changes, e.g. in management and nutrition), achieved a greater reduction of antibiotic use.

Biosecurity interventions pay off

Of course, the interventions necessary to achieve an increased level of biosecurity carry some costs. However, the interventions have proven to also improve productivity. Especially if taken with other measures such as improved management of newborn animals and nutritional improvements. The same studies which report that biosecurity improvements decrease antibiotics use also report an improvement

in animal performance. In the case of broilers, Laanen (2013) found a reduction of 0.5 percentual points in mortality and one point in FCR; and Collineau (2017) reported a reduction in mortality in pigs during both the pre-weaning and fattening period of 0.7 and 0.9 percentual points, respectively.

Execution

Although biosecurity improvements and other interventions necessary for [antibiotic reduction](#) programs are well known, continuous compliance of these interventions is often low and difficult. The implementation, application, and execution of any biosecurity program involve adopting a set of attitudes and behaviors to reduce the risk of entrance and spread of disease in all activities involving animal production or animal care. Measures should not be constraints but part of a process aimed to improve health of animals and people, and a piece of the multidisciplinary approach to reduce antibiotics and improve performance.

Designing effective biosecurity programs: consider five principles

When designing or evaluating biosecurity programs, we can identify five principles that need to be applied (Dewulf, et al., 2018). These principles set the ground for considering and evaluating biosecurity interventions:

1. Separation: Know your enemy, but don't keep it close

It is vital to have a good definition of the perimeter of the farm, a separation between high and low-risk animals, and dirty and clean internal areas on the farm. This avoids not only the entrance but the spread of disease, as possible sources of infection (e.g. animals being introduced in the herd and wild birds) cannot reach the sensitive population.

2. Reduction: Weaken your enemy, so it doesn't spread

The goal of the biosecurity measures is to keep infection pressure beneath the level which allows the natural immunity of the animals to cope with the infections (Dewulf, et al., 2018). Lowering the pressure of infection e.g. by an effective cleaning and disinfection program, by the reduction of the stocking density, and by changing footwear when entering a production house.

3. Focus: Hunt the elephant in the room, shoo the butterflies

In each production unit, some pathogens can be identified as of high economic importance due to their harm and frequency. For each of these, it is even more important, to understand the likely routes of introduction into a farm and how it can spread within it. Taking into consideration that not all disease transmission routes are equally significant, the design of the biosecurity program should focus first on high-risk pathogens and transmission routes, and only subsequently on the ones lower-risk (Dewulf, et al., 2018).

4. Repetition: When the danger is frequent, the probability of injury is increased

In addition to the probability of pathogen transmission via the different transmission routes, the frequency of occurrence of the transmission route is also highly significant when evaluating a risk (Alarcon, et al., 2013). When designing biosecurity programs, risky actions such as veterinary visits, if repeated regularly must be considered with a higher risk.

5. Scaling: In the multitude, it is easy to disguise

The risks related to disease introduction and spread are much more important in big farms (Dorea, et al., 2010); more animals may be infected and maintain the infection cycle, also large flocks/herds increase the infection pressure and increase the risk by contact with external elements such as feed, visitors, etc.

Can we still improve our biosecurity?

Almost 100% of poultry and swine operations already have a nominal biosecurity program, but not in all cases is it fully effective. BioCheck UGent, a standardized biosecurity questionnaire applied in swine and broiler farms worldwide, shows an average of 65% and 68% in conformity, respectively, from more than 3000 farms between both species (UGent, 2020). Therefore, opportunities to improve can be found in farms globally, and they pay off.

To find these opportunities, consider three situations you need to know:

1. **Know your menace:** Identify and prioritize the disease agents of greatest concern for your production system by applying the principles of **focus** and **repetition**. Consider the size of the facility when evaluating risks applying the **scaling**
2. **Know your place:** Conduct an assessment of the facility. A starting point is to define the *status quo*. For that, operation-existing questionnaires or audits can be used. However, the “new eyes principle” should be applied and an external questionnaire such as BioCheck UGent (biocheck.ugent.be) is recommended. The questionnaire will help you identify gaps in your **biosecurity plan** as well as **processes** that may be allowing pathogens to enter or move from one location to another, and measures that can be implemented applying the principles of **separation** and **reduction**.
3. **Know your processes:** Implement processes and procedures that apply the biosecurity principles and help to eliminate, prevent, or minimize the potential of disease. A deep evaluation of the daily farm processes will aid in risk mitigation, considering, among others, movement of personnel, equipment, and visitors, the entrance of pets, pests and vermin, dealing with deliveries and handling of mortality and used litter.

Compliance - The weak link in biosecurity programs

Achieving systematic compliance of biosecurity protocols on a farm is a complex, interactive, and continuous process influenced by several factors (Delabbio, 2006) and an ongoing challenge for animal production facilities (Dewulf, et al., 2018). Thus, it is clear that the biosecurity plan can only be effective if everyone on the operation follows it constantly, i.e. if everyone performs in **compliance**.

Compliance can be defined as the extent to which a person's behavior coincides with the established rules. Thus, compliance with biosecurity practices should become part of the culture of the facility. Poor

compliance in relation with biosecurity can be connected to:

- Lack of knowledge or understanding of the biosecurity protocols (Alarcon, et al., 2013; Cui & Liu, 2016; Delpont, et al., 2020)
- Lack of consequences for non-compliance (Racicot, et al., 2012a)
- A company culture of inconsistent or low application of biosecurity protocols (Dorea, et al., 2010)

In general terms, compliance with biosecurity procedures has been found to be incomplete in different studies (Delpont, et al., 2020; Dorea, et al., 2010; Gelaude, et al., 2014; Limbergen, et al., 2017). In one study (Racicot, et al., 2011) used hidden cameras, to assess biosecurity compliance in Quebec, Canada and found 44 different biosecurity fails made by 114 individuals (farm workers and visitors) in the participating poultry farms, over the course of 4 weeks; in average four mistakes were made per visit. The most frequent mistakes were ignoring the delimitation between dirty and clean areas, not changing boots, and not washing hands at the entrance of the barns; these three mistakes were committed in more than 60% of the occasions, regardless of being farm employees or visitors. These are frequent breaches not only of those farms in Quebec but found frequently in many animal production units around the world and have a high probability of causing the entrance and spread of pathogens.

How to create a high biosecurity culture: start now!

Creating, applying, and maintaining a biosecurity culture is the most effective way to make sure that compliance of the biosecurity program and procedures is high on the farm. Decreasing, therefore, the probability of entrance and spread of pathogens, reducing the use of antimicrobials, and maintaining animal health. Some actions are recommended in order to achieve a high biosecurity culture:

1. Name an accountable person

Every operation should have a biosecurity coordinator who is accountable for developing, implementing, and maintaining the biosecurity program.

This important position should be appointed having in mind that certain personality traits may facilitate performance and execution of the labor (Delabbio, 2006; Racicot, et al., 2012; Laanen, et al., 2014; Delpont, et al., 2020) such as responsibility, orientation to action, and being able to handle complexity. Additionally, expertise – years working in the industry – and orientation to learn are strategic (Racicot, et al., 2012).

2. Set the environment

When the farm layout is not facilitating biosecurity, compliance is low (Delabbio, 2006), thus the workspace should facilitate biosecurity workflows and at the same time make them hard to ignore (Racicot, et al., 2011).

3. Allow participation

It is important to mention that not only the management and the biosecurity coordinator are responsible for designing and improving biosecurity procedures. Biosecurity practices must be owned by all the farm workers and should be the social norm.

The annual or biannual revision of biosecurity measures should be done together with the farm staff. This not only serves the purpose of assessing compliance but also allows the personnel to suggest measures addressing existing -often overlooked- gaps, and to be frank about procedures that are not followed and the reasons for it. At the same time, participation increases accountability and responsibility for the biosecurity program.

4. Train for learning

Don't take knowledge for granted. Even when a person has experience in farm work and has been working in the industry for several years, their understanding and comprehension around biosecurity may have gaps.

People are more likely to do something when they see evidence of the activity's benefit. Therefore, if workers are told about the effectiveness of the practices, showing the benefits of biosecurity and analyzing the consequences of non-compliance, they are most likely to follow the procedures (Dewulf, et al., 2018). Knowledge of disease threats and symptoms also improves on-farm biosecurity (Dorea, et al., 2010), thus workers should recognize the first symptoms of disease in animals and act upon them.

Discussion of 'What if...?' scenarios to gain an understanding of the key aspects of farm biosecurity should be held on a regular basis. Workers should see examples of the benefits of compliance – and risks of noncompliance – as part of their training.

5. Lead by example

A high biosecurity culture requires everyone to comply regardless of status.

Personnel practice of biosecurity procedures is not only affected by the availability of resources and training, but also by the position that management takes on biosecurity and the feedback provided. The management and owners must transmit a message of commitment to the farm personnel, owning and following biosecurity practices, procedures and protocols, giving positive and negative feedback on the personnel's compliance, supplying information on farm performance and relating it with biosecurity compliance and ensuring adequate resources for the practice of biosecurity (Delabbio, 2006).

When necessary, management also should enforce personnel compliance by disciplinary measures, firings, and creating awareness about the consequences of disease incidence. Nevertheless, the recognition of workers' contribution to animal health performance also has a positive impact on biosecurity compliance (Dorea, et al., 2010).

The bottom line

Biosecurity is necessary for disease prevention in any animal production system. Actions and interventions that prevent the entrance and spread of disease in a production unit have a pay-off as they often lead to performance improvements and lower antimicrobial use. Maintaining a successful production unit requires a multidisciplinary approach in which [biosecurity compliance](#) needs to be taken seriously and also actions to improve in other areas such as management, health, and nutrition.

By Technical Team, EW Nutrition

References

[Alarcon, P., Wieland, B., Pereira, A. L. & Dewberry, C., 2013. Pig Farmer's perceptions, attitudes, influences, and management of information in the decision-making process for disease control. *Preventive Veterinary Medicine*, 116\(3\).](#)

[Collineau, L. et al., 2017. Profile of pig farms combining high performance and low antimicrobial usage within four European countries. *Veterinary Record*, Volume 181.](#)

[Collineau, L. et al., 2017a. Herd-specific interventions to reduce antimicrobial usage in pig production without jeopardizing technical and economic performance. *Preventive Veterinary Medicine*.](#)

Cui, B. & Liu, Z. P., 2016. Determinants of Knowledge and Biosecurity Preventive Behaviors for Highly Pathogenic Avian Influenza Risk Among Chinese Poultry Farmers. *Avian Diseases*, 60(2).

[Davies, R. & Wales, A., 2019. Antimicrobial Resistance on Farms: A Review Including Biosecurity and the Potential Role of Disinfectants in Resistance Selection. *Comprehensive Reviews in Food Science and Food Safety*.](#)

de Gussem, M. et al., 2016. *Broiler Signals*. First ed. Zutphen: Roodbont Publishers.

Delabbio, J., 2006. How farmworkers learn to use and practice biosecurity. *Journal of Extension*, 44(6).

Delpont, M. et al., 2020. Determinants of biosecurity practices in French duck farms after an H5N8 Pathogenic Avian Influenza epidemic: The effect of farmer knowledge, attitudes and personality traits. *Transboundary and Emerging Diseases*.

Dewulf, J. et al., 2018. *Biosecurity in animal production and veterinary medicine*. First ed. Den Haag: Acco Nederland.

Dorea, F., Berghaus, R., Hofacre, C. & Cole, D., 2010. Biosecurity protocols and practices adopted by growers on commercial poultry farms in Georgia USA. *Avian Diseases*, 54(3).

Gelaude, P. et al., 2014. Biocheck.UGent: A quantitative tool to measure biosecurity at broiler farms and the relationship with technical performances and antimicrobial use. *Poultry Science*.

Laanen, M. et al., 2014. Pig, cattle, and poultry farmers with a known interest in research have comparable perspectives on disease prevention and on-farm biosecurity. *Preventive Veterinary Medicine*.

Laanen, M. et al., 2013. Relationship between biosecurity and production/antimicrobial treatment characteristics in pig herds. *The Veterinary Journal*, 198(2).

Limbergen, T. et al., 2017. Scoring biosecurity in European conventional broiler production. *Poultry Science*.

Postma, M. et al., 2016. Evaluation of the relationship between the biosecurity status, production parameters, herd characteristics, and antimicrobial usage in farrow-to-finish pig production in four EU countries. *Porcine Health Management*.

Racicot, M., Venne, D., Durivage, A. & Vaillancourt, J.-P., 2011. Description of 44 biosecurity errors while entering and exiting poultry barns based on video surveillance in Quebec, Canada. *Preventive veterinary medicine*, Volume 100.

Racicot, M., Venne, D., Durivage, A. & Vaillancourt, J.-P., 2012a. Evaluation of strategies to enhance biosecurity compliance on poultry farms in Quebec: Effects of audits and cameras. *Preventive veterinary medicine*, Volume 103.

Racicot, M., Venne, D., Durivage, A. & Vaillancourt, J.-P., 2012. Evaluation of the relationship between personality traits, experience, education, and biosecurity compliance on poultry farms in Quebec, Canada. *Preventive veterinary medicine*, Volume 103.

UGent, b., 2020. *biocheck UGent*. [Online]
Available at: www.biocheck.ugent.be

Why we need to replace zinc oxide in tackling post-weaning diarrhea



Piglets experience significant stress when they are weaned from the sow and change diet, making them susceptible to gastrointestinal disorders. Primarily during the first two weeks after weaning, they are likely to suffer from post-weaning diarrhea (PWD). PWD is a significant problem for pig producers worldwide: it leads to severe dehydration, stunted growth and [mortality rates of up to 20-30%](#). Treatment and additional labor costs further squeeze farm profitability and necessitate unwanted antibiotic interventions.

Zinc oxide: an effective but highly problematic tool

Since the early 1990s zinc oxide (ZnO) has been used to control post-weaning diarrhea and promote growth in piglets, mainly at pharmacological dosages of 2500 to 3000ppm. Its mode of action is still not entirely understood; effects on [immune or metabolic processes, altered microbiota, or post-absorptive metabolism](#) are likely to play a role. What is clear is that the use of ZnO in European pig production has strongly increased since the EU banned the use of antibiotic growth promoters such as [colistin](#) in 2006 to curb the development of antimicrobial resistance.

Pigs depend on a continuous supply of zinc. Among other roles, this trace element constitutes a functional component of around 300 biochemical enzymes, making it pivotal to most metabolic processes, and by extension to optimal health, production and reproduction. [Modern pig diets thus include zinc supplementation](#) to meet the animals' requirements. The European Food Safety Authority (EFSA) currently suggests that [a total level of 150ppm](#) of zinc in feed matches the animals' physiological need for zinc. The EFSA's concerns are solely connected to the environmental concerns arising from pharmacological high dosages of ZnO.

These concerns are grave indeed: zinc is a heavy metal after all. Too much zinc is toxic for the animal, hence its physiology ensures that excessive zinc intake is excreted. The bioavailability and absorption of zinc from zinc oxide is particularly low. Therefore most of the zinc given to piglets in this way accumulates in their manure – which is widely used as an organic fertilizer for agricultural soils.

The continual application of manure gradually increases topsoil zinc concentrations; leaching and run-off then lead to contamination of groundwater, surface waters, and sediment. As zinc is non-volatile and non-degradable, it is only a matter of time before concentrations lead to ecotoxic effects, including food crops, aquatic life, and drinking water. Classic mitigation measures such as diluting the manure or keeping certain minimum distances between application areas and surface waters can only slow down the environmental accumulation of zinc, not prevent it.

EU ban: ZnO to be phased out by 2022

In 2017, the European Medicines Agency (EMA) – the EU agency responsible for the scientific evaluation, supervision and safety monitoring of medicines, including veterinary medicinal products – conducted an overall risk-benefit analysis for ZnO. It concluded that the benefits of preventing diarrhea in pigs did not outweigh the significant environmental risks caused by zinc pollution. By June 2022 [all EU member states will thus have to withdraw marketing authorizations](#) for veterinary medicinal products containing zinc oxide that are administered orally to food-producing species.

In its [decision](#), the EMA's Committee for Medicinal Products for Veterinary Use also points out the risk that, due to co-resistance, the use of zinc oxide might promote the development of antimicrobial resistance. High doses of zinc supplementation have been shown to [increase the proportion of multidrug-resistant *E. coli*](#) and [Salmonella](#), two of the most important pathogens in pig production.

What is more, studies show that [excessive zinc can accumulate in the liver, the pancreas, and blood serum](#), and that it [permanently reduces the lactobacilli population](#) of the gut flora. With what consequences for performance in the fattening phase? Hence, there are plenty of reasons why getting rid of zinc oxide is a good thing and will ultimately result in even better, more sustainable pig production – but, of course, only if effective replacement strategies to control PWD and boost piglet performance are in place.

Towards zero ZnO: smart feed additives optimize gut health

The search for ZnO alternatives takes us right back to the start, to the piglets' challenged [gastrointestinal tract](#). During their first three months of life, [pigs' gastrointestinal system undergoes a complex maturation process](#) of its epithelial, immune, and enteric nervous systems. Only once all of these systems are fully developed is the gut capable of delivering its normal functions (digestion, nutrient absorption, immunity, etc.), while also providing an effective barrier against the pathogens, antigens, and toxins in the lumen.

Unlike in nature, where weaning occurs around the time when GIT functions have matured, weaning in commercial pig production takes place during this vulnerable developmental period. Post-weaning diarrhea is ultimately a consequence of intestinal dysbiosis, a state of imbalance in the intestinal microbiome which in turn is induced by the dietary, behavioral, and environmental stressors of the weaning phase (such as separation from the sow, vaccinations, transport, introduction of solid feed).

PWD control thus starts with managing these stressors, which includes ensuring sufficient colostrum intake, gradual feed changes, and meticulous nursery hygiene. Critically, the weaning diet needs to optimally support gut health. Intelligent feed additive solutions are able to

- reduce the pathogenic load in the piglet's GIT,
- strengthen the piglet's maturing gut barrier functionality, and
- selectively induce the development of beneficial microorganisms within the microbiome.

A synergistic combination of phytomolecules, medium-chain fatty acids, glycerides of butyric acid, and prebiotics achieves these objectives in a reliable and cost-effective manner. Thanks to their antimicrobial, anti-inflammatory, and digestive properties these selected ingredients effectively support piglets during this critical phase of their postnatal gut development, while also boosting their feed intake.

In the past decade, the European pig sector has successfully adapted to the 2006 ban on antibiotic growth

promoters through significant improvements in management and feed practices. Cutting out zinc oxide is an ambitious challenge – but with the support of targeted, functional feed additives, producers will be able to set their piglets up for a strong, sustainable, zero ZnO health and growth performance.

*You can find this article in polish and italian.

References

Amezcuca, Rocio, Robert M. Friendship, Catherine E. Dewey, Carlton Gyles, and John M. Fairbrother. "Presentation of postweaning *Escherichia coli* diarrhea in southern Ontario, prevalence of hemolytic *E. coli* serogroups involved, and their antimicrobial resistance patterns." *Canadian Journal of Veterinary Research* 66, no. 2 (April 2002): 73-8. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC226986/>.

Bednorz, Carmen, Kathrin Oelgeschläger, Bianca Kinnemann, Susanne Hartmann, Konrad Neumann, Robert Pieper, Astrid Bethe, et al. "The Broader Context of Antibiotic Resistance: Zinc Feed Supplementation of Piglets Increases the Proportion of Multi-Resistant *Escherichia coli* in Vivo." *International Journal of Medical Microbiology* 303, no. 6-7 (2013): 396-403. <https://doi.org/10.1016/j.ijmm.2013.06.004>.

Brugger, Daniel, and Wilhelm M. Windisch. "Strategies and Challenges to Increase the Precision in Feeding Zinc to Monogastric Livestock." *Animal Nutrition* 3, no. 2 (March 24, 2017): 103-8. <https://doi.org/10.1016/j.aninu.2017.03.002>.

Burrough, Eric R., Carson De Mille, and Nicholas K. Gabler. "Zinc Overload in Weaned Pigs: Tissue Accumulation, Pathology, and Growth Impacts." *Journal of Veterinary Diagnostic Investigation* 31, no. 4 (June 6, 2019): 537-45. <https://doi.org/10.1177/1040638719852144>.

De Mille, Carson, Emma T. Helm, Eric R. Burrough, and Nicholas K. Gabler. "Zinc oxide does not alter ex vivo intestinal integrity or active nutrient transport in nursery pigs." Paper presented at the *Zero Zinc Summit, Copenhagen, Denmark, June 17-18, 2019*. <https://svineproduktion.dk/Services/-/media/3E0A1D2A4CAC409FAA6212B91DFEA537.ashx>.

Moeser, Adam J., Calvin S. Pohl, and Mrigendra Rajput. "Weaning Stress and Gastrointestinal Barrier Development: Implications for Lifelong Gut Health in Pigs." *Animal Nutrition* 3, no. 4 (December 2017): 313-21. <https://doi.org/10.1016/j.aninu.2017.06.003>.

Rhouma, Mohamed, Francis Beaudry, William Thériault, and Ann Letellier. "Colistin in Pig Production: Chemistry, Mechanism of Antibacterial Action, Microbial Resistance Emergence, and One Health Perspectives." *Frontiers in Microbiology* 7 (November 11, 2016): Article 1789. <https://doi.org/10.3389/fmicb.2016.01789>.

Starke, Ingo C., Robert Pieper, Konrad Neumann, Jürgen Zentek, and Wilfried Vahjen. "The Impact of High Dietary Zinc Oxide on the Development of the Intestinal Microbiota in Weaned Piglets." *FEMS Microbiology Ecology* 87, no. 2 (February 1, 2014): 416-27. <https://doi.org/10.1111/1574-6941.12233>.

Challenging times for broilers? Phytomolecules, not antibiotics, are the answer



*by **Ajay Bhoyar**, Global Technical Manager, EW Nutrition*

Anyone working with today's fast-growing broiler chicken knows that it is a sensitive creature – and so is its gut health. Thanks to continuous improvements in terms of [genetics and breeding](#), nutrition and feeding, as well as general management strategies, broiler production has tremendously upped performance and efficiency over the past decades. It is estimated that, between 1957 and 2005, the [broiler growth rate increased by over 400%, while the feed conversion ratio dropped by 50%](#).

These impressive improvements, however, have come at the cost of intense pressure on the birds' digestive system, which needs to process large quantities of feed in little time. To achieve optimal growth, a broiler's [gastrointestinal tract \(GIT\)](#) needs to be in perfect health, all the time. Unsurprisingly, enteric

diseases such as [necrotic enteritis](#), which severely damages the intestinal mucosa, hamper the intestines' capacity to absorb nutrients and induce an inflammatory immune response.

The modern broiler's gut - a high-performing, but sensitive system

However, in a system as high performing as the modern broiler's GIT, much less can lead to problems. From when they are day-old chicks up to slaughter, broilers go through several challenging phases during which they are more likely to show impaired gut functionality, e.g. after vaccinations or feed changes. [Good management practices go a long way towards eliminating unnecessary stressors](#) for the animals, but some challenging periods are unavoidable.

The transition from starter to grower diets is a classic situation when nutrients are very likely to not be well digested and build up in the gut, fueling the proliferation of harmful microbes. Immunosuppressive stress in combination with an immature intestinal microflora results in disturbances to the bacterial microbiota. At "best", this entails temporarily reduce nutrient absorption, in the worst case the birds will suffer serious intestinal diseases.

Phytomolecules - the intelligent alternative to antibiotics

To safeguard performance during stressful periods, poultry producers need to anticipate them and proactively provide effective gut health support. For many years, this support came in the form of antibiotic growth promoters (AGP): administered prophylactically, they were effective at keeping harmful enteric bacteria in check. However, due to grave concerns about the [development of antimicrobial resistance](#), non-therapeutic antibiotics use has been banned in many countries. Alternatives need to focus on improving feed digestibility and strengthening gut health, attacking the root causes of why the intestinal microflora would become unbalanced in the first place.

Phytomolecules are secondary metabolites active in the defense mechanisms of plants. Studies have found that certain phytomolecules [stimulate digestive enzyme activities](#) and stabilize the gut microflora, "leading to improved feed utilization and less exposure to growth-depressing disorders associated with digestion and metabolism" (Zhai et al., 2018). With other trials showing [positive effects on broilers' growth performance and feed conversion](#), the research indicates that phytomolecules might also specifically support chickens during challenging phases.


The effect of phytomolecules on broilers during a challenging phase

A study was conducted over a period of 49 days on a commercial broiler farm of an AGP-free integration operation in Japan. The farm reported gut health challenges in the second and third week of the fattening period due to vaccinations and changes to the animals' diets. The trial included 15504 Ross 308 broilers, divided into two groups. The negative control group included a total of 7242 birds, kept in another house.

All the birds were fed the standard feed of the farm. The trial group (8262 birds) received Activo Liquid, which contains a synergistic combination of phytomolecules, administered directly through the drinking water. Activo Liquid was given at an inclusion rate of 200ml per 1000L of water (3.3 US fl oz per gallon of stock solution, diluted at 1:128), from day 8 until day 25, for 8 hours a day.

The results are summarized in Figure 1:

Figure 1: Improved broiler performance for Activo Liquid group (day 49)



The Activo Liquid group clearly showed performance improvements compared to the control group. Livability augmented by 1.5%, while the feed conversion rate improved by 3.2%. This resulted in a more than 5% higher score in terms of the performance index.

Challenging times? Tackle them using phytomolecules

Poultry producers take great care to eliminate unnecessary sources of stress for their birds. Nonetheless, during their lifecycle, broiler chickens face challenging periods during which the balance of the intestinal microflora can easily become disturbed, with consequences ranging from decreased nutrient absorption to full-blown enteric disease.

The trial reviewed here showed that, after receiving Activo Liquid, broilers raised without AGPs showed encouraging performance improvements during a challenging phase of feed changes and vaccinations. Likely thanks to the activation of digestive enzymes and a stabilization of the gut flora, the broilers showed improved livability and feed conversion, thus delivering a much more robust performance during a critical phase of their lives. In times where the non-therapeutic use of antibiotics is no longer an option, phytomolecules allow poultry farmers to effectively support their animals during challenging times.

References

Photo Source: [Aviagen](#)

[Adedokun, Sunday A., and Opeyemi C. Olojede. "Optimizing Gastrointestinal Integrity in Poultry: The Role of Nutrients and Feed Additives." *Frontiers in Veterinary Science* 5 \(January 31, 2019\): 348.](#)

[Jamroz, D., T. Wiertelicki, M. Houszka, and C. Kamel. "Influence of Diet Type on the Inclusion of Plant Origin Active Substances on Morphological and Histochemical Characteristics of the Stomach and Jejunum Walls in Chicken." *Journal of Animal Physiology and Animal Nutrition* 90, no. 5-6 \(March 23, 2006\): 255-68.](#)

[Tavárez, Marcos A., and Fausto Solis De Los Santos. "Impact of Genetics and Breeding on Broiler Production Performance: a Look into the Past, Present, and Future of the Industry." *Animal Frontiers* 6, no. 4 \(October 1, 2016\): 37-41.](#)

[Zhai, Hengxiao, Hong Liu, Shikui Wang, Jinlong Wu, and Anna-Maria Kluefer. "Potential of Essential Oils for Poultry and Pigs." *Animal Nutrition* 4, no. 2 \(June 2018\): 179-86.](#)

[Zuidhof, M. J., B. L. Schneider, V. L. Carney, D. R. Korver, and F. E. Robinson. "Growth, Efficiency, and Yield of Commercial Broilers from 1957, 1978, and 2005." *Poultry Science* 93, no. 12 \(December 2014\): 2970-82.](#)

Beyond AGPs: Controlling necrotic enteritis through gut health optimization



Antibiotic growth promoters (AGPs) have routinely been used in intensive poultry production for improving birds' performance. However, in recent years, reducing the use of [antibiotics in animal production has become a top priority](#), due to concerns about the development of antibiotic-resistant bacteria and mounting consumer pressure. Multiple countries have introduced bans or severe restrictions on the non-

therapeutic use of antibiotics, including in the US, where the Food and Drug Administration has implemented measures to curb the use of antibiotics since 2017.

However, the removal of AGPs poses challenges for poultry performance, including reduced feed efficiency, decreased daily weight gain, as well as higher mortality. Moreover, the withdrawal of AGPs in feed is widely recognized as one of the predisposing factors for necrotic enteritis (NE). NE is one of the most common and economically important poultry diseases, with an [estimated global impact of US\\$ 5 to 6 billion per year](#). As a result of withdrawing AGPs, the usage of therapeutic antibiotics to treat NE has increased. To break out of this vicious cycle and to secure the efficiency of poultry production, alternatives are needed that combat NE where it starts: in the gut.

Necrotic enteritis: a complex disease

NE is caused by pathogenic strains of *Clostridium perfringens* (CP): ubiquitous, gram-positive, spore-forming anaerobic bacteria. The spores of CP can be found in poultry litter, feces, soil, dust, and contaminated feed. Low levels of different CP strains are naturally present in the intestines of healthy birds, kept in check by a balanced microbiome. However, when gut health is compromised, [pathogenic strains can proliferate at the expense of unproblematic strains](#), resulting in clinical or sub-clinical NE.

Animals suffering from the clinical form show symptoms such as general depression, reluctance to move, and diarrhea, with mortality rates of up to 50%. Infected birds suffer from degenerated mucosa lesions in the small intestines. Even in its “mild”, subclinical form, which often goes unnoticed, the damage to the animals’ intestinal mucosa can result in permanently reduced performance and consequent economic losses for the producer.

Certain [predisposing factors](#) have been found to enable the proliferation of pathogenic strains in the gastrointestinal tract. Diet is a key example: the composition of the gut flora is directly linked to feed composition. High inclusion rates of cereals (barley, rye, oats, and wheat) that contain high levels of non-starch polysaccharides (NSPs), high levels of indigestible protein, and inclusion of proteins of animal origin (e.g. fishmeal) have been shown to predispose birds to NE.

A range of diseases (e.g. chicken infectious anemia, Gumboro, and Marek’s disease), but also other factors that have immunosuppressive effects, such as heat or cold stress, [mycotoxins](#), feed changes, or high stocking density, render birds more susceptible to intestinal infections. The single most prominent predisposing factor for the occurrence of NE is the [mucosal damage caused by coccidiosis](#).

Gut health is key to combating necrotic enteritis

To control NE, a holistic approach to optimizing the intestinal health of poultry is needed. It should take into account not only parameters such as diet, hygiene, and stress, but should also make use of innovative tools.

Phytomolecules, also known as secondary plant compounds, are essentially plants’ defense mechanisms against pathogens such as moulds, yeasts, and bacteria. [Studies have demonstrated the antimicrobial effects](#) of certain phytomolecules, including against antibiotic-resistant pathogens. Phytomolecules have also been found to boost the production of digestive enzymes, to suppress pro-inflammatory prostaglandins and have antioxidant properties. These features make them a potent tool for optimizing gut health, potentially to the point of replacing AGPs.

Can phytomolecules mitigate the impact of necrotic enteritis?

To study the impact of phytomolecules on the performance of broilers challenged with a NE-causing CP strain, a trial was conducted at a US-based research facility. In this 42-day study, 1050 male day-old Cobb 500 broiler chicks were divided into 3 groups, with 7 replicates of 50 chicks each.

On the first day, all animals were vaccinated against coccidiosis through a live oocyst spray vaccination. The experimental diets met or exceeded the National Research Council requirements, and were fed as crumbles/pellets. On days 19, 20, and 21, all pens, except the negative control group, were challenged with a broth culture of *C. perfringens*. A field isolate of CP known to cause NE (originating from a commercial broiler operation) was utilized as the challenge organism. On day 21, three birds from each pen were selected, sacrificed, group weighed, and examined for the degree of present NE lesions.

The positive control group received no supplements. The trial group received a synergistic combination of two phytogenic products containing standardized amounts of selected, microencapsulated phytomolecules: an in-feed phytogenic premix (Activo, EW Nutrition GmbH) and a liquid complementary feed supplied via the drinking water (Activo Liquid, EW Nutrition GmbH). The products were given at inclusion rates corresponding to the manufacturer’s baseline antibiotic reduction program recommendations (Figure 1):

Figure 1: Trial design

Trial Groups	Challenge with NE-causing CP strain	Supplements
Negative control	No	No
Positive control	Yes	No
Activo + Activo Liquid	Yes	Activo 100g/MT + Activo Liquid at 250ml per 1000l on days 12-14 for 24 hrs per day, on days 19-21 for 16 hrs per day

The trial results indicate that the addition of phytomolecules helps to mitigate the impact of NE on broilers’ performance. The group receiving Activo and Activo Liquid showed a better feed conversion (Figure 2) compared to the positive control group (NE challenge, no supplement). Also, better lesion scores were noted for animals receiving phytomolecules (0.7 and 1) than for the positive control group (1.6).

The most significant effect was observed concerning mortality: the group receiving Activo and Activo Liquid showed a 50% lower mortality rate than the positive control group (Figure 3). These results clearly indicate that phytomolecules can play an important role in mitigating losses due to NE.

Figure 2: Adjusted FCR





Figure 3: Lesion scores and mortality



Tackling necrotic enteritis in a sustainable way

In an age of AGP-free poultry production, a concerted focus on fostering animals' gut health is key to achieving optimal performance. This study strongly demonstrates that, thanks to their antimicrobial, digestive, anti-inflammatory and antioxidant properties, phytomolecules effectively support birds' intestinal health when challenged with NE. The inclusion of [Activo](#) and Activo Liquid, two phytogenic products designed to synergistically support birds during critical periods, resulted in improved feed conversion, better lesion scores, and 50% lower mortality.

In combination with good dietary, hygiene, and management practices, phytomolecules are therefore a potent tool for reducing the use of antibiotics: including Activo and Activo Liquid in their animals' diets allows poultry producers to reduce the incidence of NE, to mitigate its economic impact in case of outbreaks, and therefore to control NE in a sustainable way.

By by Ajay Bhoyar, Global Technical Manager, EW Nutrition

References

[Antonissen, Gunther, Siska Croubels, Frank Pasmans, Richard Ducatelle, Venessa Eeckhaut, Mathias Devreese, Marc Verlinden, Freddy Haesebrouck, Mia Eeckhout, Sarah De Saeger, Birgit Antlinger, Barbara Novak, An Martel, and Filip Van Immerseel. "Fumonisin Affects the Intestinal Microbial Homeostasis in Broiler Chickens, Predisposing to Necrotic Enteritis." *Veterinary Research* 46, no. 1 \(September 23, 2015\): Article 98. doi:10.1186/s13567-015-0234-8.](#)

[Moore, Robert J. "Necrotic Enteritis Predisposing Factors in Broiler Chickens." *Avian Pathology* 45, no. 3 \(May 31, 2016\): 275-81. doi:10.1080/03079457.2016.1150587.](#)

Tang, Karen L., Niamh P. Caffrey, Diego B. Nóbrega, Susan C. Cork, Paul E. Ronksley, Herman W. Barkema, Alicia J. Polachek, Heather Ganshorn, Nishan Sharma, James D. Kellner, and William A. Ghali. "Restricting the Use of Antibiotics in Food-producing Animals and Its Associations with Antibiotic Resistance in Food-producing Animals and Human Beings: A Systematic Review and Meta-analysis." *The Lancet Planetary Health* 1, no. 8 (November 6, 2017): 316-27. doi:10.1016/s2542-5196(17)30141-9.

Van Immerseel, Filip, Julian I. Rood, Robert J. Moore, and Richard W. Titball. "Rethinking Our Understanding of the Pathogenesis of Necrotic Enteritis in Chickens." *Trends in Microbiology* 17, no. 1 (2009): 32-36. doi:10.1016/j.tim.2008.09.005.

[Wade, Ben, and Anthony Keyburn. "The True Cost of Necrotic Enteritis." *PoultryWorld*. October 09, 2015. Accessed August 19, 2019.](#)

Source Photo: [Aviagen](#)