Feed and water management strategies to mitigate heat stress in layers



Dr Daniel Valbuena, Global Manager of Technical Services, Hy-Line International - Conference Report

Feed and water management strategies are essential to help mitigate the negative effects of heat stress on bird welfare, production, and profitability. In EW Nutrition's Poultry Academy in September, the topic was approached in a comprehensive and practical presentation.

Feed management

Feed consumption of the flock should be closely monitored during hot weather. It is important to rebalance the diet for critical nutrients, particularly amino acids, calcium, sodium and phosphorous according to the birds' productivity demand (i.e., stage of production) and the observed feed intake. Insufficient amino acid intake is the primary reason for productivity loss during hot weather. Several strategies may be employed to help to manage elevated temperatures and maintain higher levels of feed intake:

- Withdrawing feed from birds 6 hours before peak hot temperatures in the afternoon can lower the risk of heat stress. Encourage as much consumption as possible in the early morning or evening. Using lighting for midnight feeding encourages feed intake.
- One third of the daily feed ration should be given in the morning and two thirds in the late afternoon. An additional advantage is the availability of calcium in the digestive system during shell formation at night and in the early hours of the morning. This will improve shell quality and reduce the birds from depleting bone calcium.
- Normally a maximum 1 hour for feeder clean-out time is recommended, but this can be

extended to 3 hours when the temperature exceeds 36°C.

- Consider adding a 1-2-hour midnight feeding.
- Alter feed particle size, either by increasing it or by feeding a crumble diet. With crumble diets in laying flocks, a supplementary source or presentation of large particle limestone is recommended.
- Formulate diets using highly digestible materials, particularly protein sources. Metabolism of excess protein is particularly heat-loading on the bird. Formulate to digestible amino acid targets and do not apply a high crude protein minimum in the formula. Synthetic amino acids can reduce crude protein in the diet without limiting amino acid levels.
- Increasing the proportion of energy contribution from highly digestible lipid, rather than starches
 or proteins, will reduce the body heat production resulting from digestion. This is known as heat
 increment and is lowest with the digestion of dietary fat.
- The bird's metabolizable energy requirement decreases as ambient temperature increases to above 21°C, resulting from a reduction of energy requirements for maintenance. The energy requirement will decrease with the rise of temperature up to 27°C, above which it will start to increase again since the bird needs additional energy for panting to reduce body heat.



Management schedule during times of heat stress

Water management

During periods of high environmental temperature, birds have a high demand for drinking water. The water-to-feed consumption ratio is normally 2:1 at 21°C but increases to 8:1 at 38°C. Adequate drinking water must be available to heat-stressed flocks. Ensure that drinkers have sufficient water flow (>70 mL/minute/nipple drinker). If water flow is less the lines need to be checked for flow restriction. If there's a build-up of iron and other minerals, it needs to be removed. Don't forget to routinely check water filters and replace them as needed.

It's easy to overlook a non-functioning drinker here and there; drinkers must be systematically checked to

make sure they're all working. For floor-reared flocks, providing additional drinkers can help accommodate the increased water consumption.

During hot weather, you need to ensure your water system can accommodate the bird's increased water consumption, <u>and</u> the additional water demands for foggers, evaporative cooling systems and roof sprinklers. The availability of drinking water to a heat-stressed flock should never be compromised.

Cool water temperatures (<25°C) will encourage the birds to drink and reduces the birds' core temperature. Flush water lines and waterers routinely to keep the water fresh and cool, increasing water consumption, and sustaining egg production. If available, ice can also be added to header tanks. When mechanical cooling systems fail, water flushing can serve as an emergency measure during heat stress.

Drinking water from overhead water tanks can become hot if exposed to direct sunlight. These water tanks should be a light color, insulated and covered to avoid direct sunlight. Water tanks are ideally placed inside the house or underground. Water pipes in the house should not be installed close to the roof to avoid heat from the roof warming up the water in the pipes.





Having the water tank inside the house (above) or light-colored and covered to avoid direct sunlight (below) keeps the water cooler

Use vitamin (A, D, E and B complex) and electrolyte supplements in the drinking water to replenish the

loss of sodium, chloride, potassium, and bicarbonate in the urine. Electrolyte supplements are best used in anticipation of a heat stress period and can be added to drinking water for up to 3 days.

Coping with evolutions in the performance and nutritional requirements of layers



Dr. Vitor Arantes, Global Technical Services Manager and Global Nutritionist, Hy-Line International – Conference Report

The layer industry has gone through significant changes during the past decades and has a remarkable capacity to cope with new challenges. Dr Vitor Arantes, Global Technical Services Manager and Global Nutritionist, Hy-Line International, noted that increased egg production, improved feed efficiency, and adaptation of egg quality and bird welfare to consumer preferences have contributed significantly to the success of the egg industry. However, continuous improvement in egg production per hen housed is the most important selection criteria in layer breeding.

Egg producers needs include:

- More saleable eggs,
- Eggshell quality,
- Easier behaviour
- Housing systems
- Egg size specifics
- Sanitary / environmental challenges
- Profits through productivity

Primary breeders can deliver these producer needs through:

- Having the correct product for each country
- Constant follow up
- Local presence, trust relationship
- Accurate data collection
- Critical data analysis
- Understand the company's goals
- Customized technical services according to each customer needs

How has genetics changed?

Examples of genetic progress in layers from 1984 to 2022 cited by Dr Arantes include:

- Higher persistency (+30 weeks >90%)
- Higher egg mass (+5.5 kg/hen housed)
- Smaller hen (-21% mature body weight)

Dr Arantes states the record clutch size, defined as the unstopped length of individual egg production on a daily basis, was an amazing 474 days for a White Plymouth Rock hen. This genetic progress necessitates adjustments in nutrition and management.



As shown below, growth and organ development occur at various ages. "There is no margin for mistakes – a lack of growth during a stage <u>could have a detrimental impact on pullet quality and subsequent</u> <u>production</u>," stressed Dr Arantes.

Multi-phasic growth and development during rearing and start of lay

System	Age (weeks)	Consequence	
Gastrointestinal	0-6	Shorter intestinal tract/reduced nutrient absorption	
Immune	0-6	Flocks more susceptible to disease challenges	
Skeleton	6-12	Shorter frames/less calcium reserves	
Muscle	6-12	Impact in persistency of production	
Fat	>12	Excess can lead to fatty liver, prone to prolapse and mortality	

0-6 weeks of age

Most of the development of the organs of the digestive tract and the immune system occurs during the first 6 weeks of age. Problems that occur during this period can have negative effects on the function of these systems. Birds stressed during this period may have lifelong difficulties in digesting and absorbing feed nutrients. Immunosuppression may also result from problems during this period, leaving the bird more susceptible to diseases and less responsive to vaccinations.

6-12 weeks of age

Most of the adult structural components – muscles, bones and feathers are obtained during the period of rapid growth that occurs at 6-12 weeks of age. Growth deficiencies during this period will prevent the bird from obtaining sufficient bone and muscle reserves, which are necessary to sustain a high level of egg production and to maintain good eggshell quality. About 95% of the skeleton is developed at the end of the bird's 13 weeks of life. At this time, the plates of the long bones become calcified and further growth in bone size cannot occur.

12-18 weeks of age

During this period, the growth rate slows, and the reproductive tract matures and prepares for egg production. Muscle development continues and the proliferation of fat cells takes place. Excessive weight gain during this period can result in an excessive amount of abdominal fat. Low body weight and stressful events at this time can delay the start of egg production. From 7-10 days before oviposition of the first egg, the medullary bone that is located within the cavities of the long bones can be increased by feeding the bird a pre-laying ration with higher levels of calcium than the development stage.

Bodyweight is a key factor for flock management as this will influence future performance of birds. Consequently, bodyweight should be controlled during the whole life of the layer flocks. Management, in particular nutrition and lighting programs, can help to control bodyweight so birds can achieve their genetic potential.

Uniformity

Uniformity is the most important KPI in our business. However, with the trend towards larger flocks, maintaining uniformity is becoming more challenging. With larger flocks, it is difficult to source one unique flock which thus usually comprises multiple breeding flocks of different ages. Inevitably, uniformity will be poor, hence the need for tools to address unexpected issues. Lack of uniformity becomes a self-perpetuating cycle – dominant versus dominated.

Many egg producers use average body weights compared to the breeder recommendations as a guide to

flock status. However, knowing if you have good body weight uniformity is another valuable management tool. In any flock some birds are lighter or heavier than the average body weight. Poor uniformity makes management decisions, such as lighting, feed amounts or diet phase more difficult.

Ideally, the body weight coefficient of variation (CV) should be +/-10% of the mean, increasing the likelihood that your management decision will be appropriate for most of the flock. Inappropriate diet changes, bird handling, vaccination and transfer can reduce uniformity. Flocks should be at 90% uniformity at the time of transfer to the laying facility. Body weight at point of lay significantly affected egg production and eggshell quality.

Grading into 2 or 3 sub-populations of different average bodyweights may be necessary so that each group can be managed in a way that will achieve good whole flock uniformity at the point of lay. The best predictor of future laying performance is the pullet's body weight and body type at the point of lay.

Vision egg

Vision Egg is a custom diagnostic tool used to analyze data and emphasize flock performance to achieve the highest genetic potential from Hy-Line layers with recommendations connected to customer profitability. This growing, robust database includes data from over 1 billion hens strengthens our flock performance diagnostic tool for improved profitability for Hy-Line customers.

<u>Hy-Line</u> customers can take advantage of this opportunity by sending flock data to their regional business manager or technical service specialist. The information shared with Hy-Line is kept completely confidential.

Summary

The challenge is not egg numbers, stated Dr. Arantes, but saleable eggs. Correct body weight and high uniformity of the flock at point of lay will result in good performance over the laying period, with high peak production and good persistency of production and the production of good quality eggs. Management is the key factor to regulation of body weight during rearing and at point of lay.

How to mitigate formulation costs when ingredient prices are high



Conference Report

The price of corn and soybeans dictates the price of all other ingredients, including to some extent amino acids, stated Dr Steve Leeson Professor Emeritus, University of Guelph, Canada at the recent <u>EW Nutrition</u> Poultry Academy in Jakarta, Indonesia.

The big question is, when times get tough, can we reduce safety margins and still get good performance?, asked Dr Leeson. "When we formulate diets, we build in some insurance. But so do the breeding companies in their recommendations. For sure, reducing safety margins takes us out of our comfort zones, but we need to be nutritionists, not mathematicians," he stressed.

Protein and energy are now expensive. As a result of this economic pressure, there is a focus on strategies to reduce feed costs and improving the production efficiency and profitability of poultry enterprises. Feed cost/kg body weight gain is not always at the lowest feed:gain.

To help achieve these targets, Dr Leeson discussed feeding and management strategies that take into account the cost mitigation requirement.

Optimize current digestibility/efficiency

With high feed prices, it is especially important to review the use of feed additives that optimize nutrient release and improve 'digestibility'. The most obvious class of such additives are the various exogenous enzymes that improve the availability of phosphorus, energy, and amino acids. In most instances, these different classes of enzymes are additive in terms of nutrient release, since they have different target substrates or modes of action. All too often, the position is taken that "I take energy uplift from my amylase, so I can't expect energy release from phytase or protease".

The energy release from phytase is invariably net energy related to removal of the phytate molecule, which in effect is an 'antigen' and takes energy to counter its negative effects. The energy release from an amylase, however, is obviously related simply to the improved digestibility of carbohydrate complexes. Similarly, a protease enzyme will always provide energy, since all protein/amino acids are eventually used for energy during protein turnover, hence our use of the often forgotten 'n' in AMEn. We also have the choice of enzyme concentration, especially for phytase, which in the current economic solution is likely to be close to 2 – 2.5 doses, assuming a single dose is around 500-600 FTUs. The economics of super-dosing or mega-dosing is greatly impacted by the cost of the enzyme.

The response of phytase varies with individual amino acids, and with ingredients, with greater responses with ingredients of lower inherent digestibility. Generally, Dr Leeson suggests that a protease will capture 20% of indigestible amino acids. For example:

- 70% digestibility = +6% uplift
- 90% digestibility = +2% uplift

Relax ingredient constraint maximums

Probably the greatest current cost savings can be made from relaxing the maximum levels on ingredients. While corn and soybean meal levels are usually without restriction, we often impose limits on the upper levels of 'alternative' ingredients such as distillers grains, rice by-products and rapeseed/canola meals, etc. When the upper levels are reached in the formula, this suggests cost savings from using higher levels. Current restraints are based on past knowledge of perhaps variable nutrient composition and so the decision to use more of any ingredient must be based on past knowledge of on-going quality control assays. Although we can achieve considerable detail today in such QC assays, monitoring for (consistency of) crude fiber, crude protein, fat, and moisture alone, provide a sound basis for decisions on whether to use more of an individual ingredient.

Source alternate ingredients

Another option is to consider 'new' alternative ingredients. In reality, however, there are no new ingredients as such, since all monogastric nutritionists around the world have only around 19 ingredients available in sufficient quantities to sustain large-scale modern feed mills. There are certainly smaller quantities of specialised local by-products that can be used to advantage, yet these are becoming scarce. Therefore, an ingredient is only novel to you, since inevitably the same ingredient has been used for many years in other regions. As such, there is a wealth of information available on the nutritive value of these 'new' ingredients that can be simply transposed to our formulation matrices.

The bird is very adaptable to new ingredients, in fact it is more responsive to nutrients. Unless there are toxins, antinutritional factors, or other negative factors, it doesn't matter to the bird. Knowing the ingredient composition is the critical feature regarding the success or failure with new ingredients.

Reduce nutrient density

Both layers and meat birds still eat quite precisely to their energy requirements. They are amazingly adaptable to a vast range of nutrient densities, assuming that they can eat enough feed as the lower levels of feed energy are approached. Success in using lower levels of nutrient density is invariably negatively impacted by factors such as high stocking density and a high environmental temperature. Conversely, reducing diet energy usually has the hidden advantage of improved pellet quality.

The key to successful use of lower energy diets lies in prediction of change in feed intake and corresponding adjustment to all other nutrients in the diet.

Flexible cost of Dietary electrolyte balance (DEB)

When first introduced in the 1970s, maintaining DEB around 250MEq was seen to optimize broiler performance, especially leg condition. There is now less emphasis on this, perhaps because of genetic selection for skeletal integrity. DEB, however, may be important during heat stress to stimulate water intake and control manure moisture. Formulating to fixed DEB levels always adds costs. Instead, Dr Leeson suggested to focus on sodium and chloride at a ratio of 1:1.3.

Optimize feed texture (pelleting)

The first consideration is to make a good quality pellet, then worry about pellet size, noted Dr Leeson. He also added he was "a big fan of sunflower meal – it's great for pellet quality."

When given a choice in particle sizes, birds invariably show a preference for the largest particles. This situation becomes obvious when 'fines' accumulate in the feeder pans over time. As shown below, as pellet size increases, so does the bird's need to consume fewer pellets. As a result, they need to spend less time at the feeder. Naturally, this idealised pellet size must be balanced against the willingness of mill managers to accommodate the necessary changes in pellet die size. Matching pellet size to bird age becomes critical as stocking density increases.

Impact of pellet size on pellet number consumed by a 30-day-old broiler

Pellet size (diameter)	4 mm length	6 mm length
3 mm	580	390
4 mm	330	220
5 mm	210	140

In the end, cost mitigation should not require complex mathematics. Nutritionists should be able to play with several types of improvements without affecting health and performance.

<u>EW Nutrition</u>'s Poultry Academy took place in Jakarta and Manila in early September 2023. Dr. Steve Leeson, an expert in Poultry Nutrition & Production with nearly 50 years' experience in the industry, was the distinguished keynote speaker.

Dr. Leeson had his Ph.D. in Poultry Nutrition in 1974 from the University of Nottingham. Over a span of 38 years, he was a Professor in the Department of Animal &Poultry Science at the University of Guelph, Canada. Since 2014, he has been Professor Emeritus at the same University. As an eminent author, he has more than 400 papers in refereed journals and 6 books on various aspects of Poultry Nutrition & Management. He also won the American Feed Manufacturer's Association Nutrition Research Award (1981), the Canadian Society of Animal Science Fellowship Award (2001), and Novus Lifetime Achievement Award in Poultry Nutrition (2011).

Metabolic disorders and muscle defects



Conference Report

At the recent <u>EW Nutrition</u> Poultry Academy in Jakarta Indonesia, Dr Steve Leeson, Professor Emeritus, University of Guelph, Canada, defined metabolic disorders as: non-infectious, occurring with adequate diets in 'normal' conditions, and mostly species-specific. Their incidence is negatively correlated to productivity. Although they often have a major genetic component, genetic selection to manage the problem is often a last resort, as there is usually a negative correlation with productivity.

Ascites

First reported in the 1970s, ascites or 'water belly' is probably the number one metabolic issue today. It is the accumulation of fluid in the abdomen, which is caused by a cascade of events related to the need to supply high levels of oxygen to the tissues. The condition was initially most prevalent in fast-growing male broilers maintained at high altitude and where there is a degree of cold stress, but nowadays the problem can occur at any altitude. In extreme situations up to 8% mortality is seen, although 1-3% mortality is currently more common. The disorder is now re-emerging with faster growth rates, as growth rate is easily the main contributing factor.

Options to limit ascites include:

- Limit growth rate
- Feed texture (mash vs. pellets)
- Never let the temperature get below 15°C for any age of bird
- Brooding ventilation economics of air flow vs. temperature
- Minimize environmental contaminants, such as dust
- Lighting programs (4-6 hours of darkness)

Sudden death syndrome (SDS)

SDS almost always affects males birds close to market weight. It frequently afflicts 1-5% of the flock and from 21-35 days it will usually be the major cause of death. Afflicted birds appear healthy, are well fleshed and invariably have feed in their digestive tract. Death occurs within 1-2 minutes, the birds most frequently being found dead on their backs. There are few changes in gross pathology. The heart may contain blood clots, that are likely post-mortem in origin, and the ventricles are usually empty. Diagnosis is usually by exclusion of other diseases. The lungs are often oedematous, although this usually occurs when birds spend time on their backs and fluid drains to the lung region by gravity. There are no specific changes in the tissue or blood profile that can be used for diagnosis. The condition is precipitated by fast growth rate, and so conversely it can be prevented by varying degrees of nutrient restriction.

Spiking mortality syndrome (SMS)

SMS is characterized by severe unexplained hypoglycemia, and always occurs from 18-21 days of age. There are few post-mortem observations, so it is often misdiagnosed. Mortality can be 2-3%. Males are more susceptible than females, probably because they are growing faster. Birds fed all-vegetable diets may be more prone to SMS. Supplementing an all-vegetable diet with milk-powder (which is high in serine), casein or serine is recommended and results in increased blood glucose.

Skeletal integrity

This disorder is not due to increased bodyweight of broilers, as the broiler is capable of supporting weight that far exceeds its own body weight. Instead, it's due to shifting the bird's center of gravity forward as breast muscle yields have increased, moving the legs further apart which puts torsional pressure on the head of the femur. Not only does it cause on-farm problems, but also complications with mechanical processing.

Imbalanced nutrient supply, such as excess of chloride, or infection with bacteria, viruses, and particularly mycoplasmas are involved.

Tibial dyschondroplasia (TD)

TD is due to abnormal cartilage development. Failure of normal vascularization limits mineralization. TD is characterized by enlargement of the hock, twisted metatarsi, and slipped tendons. A low electrolyte balance (<200MEq), high chloride (>0.3%), or low Ca:P or high P:Ca can precipitate TD. Adding manganese and choline to the diet will largely eliminate it.

Perosis

Now often termed Chondrodystrophy, it has manganese or choline deficiency as the classical cause, but it can also be seen with other B-vitamin deficiencies. As with TD, it can be aggravated by some grain fumigants.

Kinky back

Also known as Spondylolisthesis, it is not really a metabolic disorder, as *Enterococcus* infection is the most common cause. Chickens with kinky back syndrome are often seen sitting on their tail, extending their feet outward or letting them fall over to one side of their body. Once the condition stops birds from being able to walk, they are unable to reach food or water on their own and are at risk of dying from starvation. There is no treatment for kinky back.

Gizzard erosion and proventriculus

Although gizzard lesions are very common, Dr Leeson suspects their importance is overemphasized. Gizzard condition is seen in both layer and broiler chickens, but the incidence is more in broilers.

Access to grit and inclusion of at least 20% cereal particles larger than 1 mm in size in the diet will have a positive effect on the development and functioning of the gizzard and it will also reduce the frequency and severity of gizzard lesions in poultry. Ingestion of non-soluble fibers has been shown to exert strong effects on the structure and function of the gizzard. Inclusion of at least 3% coarse fibers in the feed increased the relative weight of the gizzard and reduced the pH of the gizzard contents suggesting a preventive effect of fiber.

Proventriculus appears as a very large organ and is often associated with gizzard erosion. When the proventriculus glands are affected, there is a lower secretion of hydrochloric acid and enzymes and therefore more undigested feed arrives to the intestine, where it can act as a substrate of pathogens and start digestive infections.

Breast muscle defects

Breast muscle defects are not problematic for the bird, efficiency/economics of growth, or a food safety issue. The main issue is seen at primary or secondary processing, and consumer acceptance. Due to the fast muscle growth and the enlarged muscle cells, the space between muscle fibers is reduced. This restricts the blood supply to the muscles, which can no longer reach the desired oxygen levels.

White-striping

White striping is a quality factor in chicken breast meat caused by deposits of fat in the muscle during the bird's growth and development. It is like marbling in red meat. Dr Leeson joked that it be promoted as marbled chicken – like Wagyu beef. Because hypoxia is associated with white striping, it was thought that arginine supplementation could help with vasodilation, thus supplying the muscles with better oxygen resources.

Wooden breast (WB)

WB is an <u>emerging quality defect</u>. Macroscopically, it is characterized by palpably hard, pale ridge-like bulges at the caudal end, along with clear viscous fluid, small hemorrhages, and white striping, that may occur separately or together. The main cause is the high growth rate and high breast meat yield. There is no nutritional or management solution.

Wooden breast is common in male broilers >2.5 kg bodyweight, and the incidence tends to increase with the size of the breast fillet. As the incidence of wooden breast increases, the incidence of white striping tends to decrease. Due to the visual defects and hard and chewy texture, consumers have a low acceptance of WB fillets, and they are usually downgraded to use for ground products.

Reducing oxidative stress and supplying more oxygen to the cells, enabling the muscle cells to grow very fast without meat loss will reduce the incidence of WB.

<u>EW Nutrition</u>'s Poultry Academy took place in Jakarta and Manila in early September 2023. Dr. Steve Leeson, an expert in Poultry Nutrition & Production with nearly 50 years' experience in the industry, was the distinguished keynote speaker.

Dr. Leeson had his Ph.D. in Poultry Nutrition in 1974 from the University of Nottingham. Over a span of 38 years, he was a Professor in the Department of Animal &Poultry Science at the University of Guelph, Canada. Since 2014, he has been Professor Emeritus at the same University. As an eminent author, he has more than 400 papers in refereed journals and 6 books on various aspects of Poultry Nutrition &

Management. He also won the American Feed Manufacturer's Association Nutrition Research Award (1981), the Canadian Society of Animal Science Fellowship Award (2001), and Novus Lifetime Achievement Award in Poultry Nutrition (2011).

Meat quality is a result of genetics, feeding, the microbiome, and the handling of animals and meat



by Dr. Inge Heinzl, Editor EW Nutrition

Nowadays, nutrition is no longer about pure nutrient intake; enjoyment is also a priority. Consumers attach great importance to the high quality of food and, therefore, also of meat. The genetic selection for faster growth and feeding high-energy diets made meat production more efficient and shortened the raising period. However, this selection may sometimes also result in challenges to meat quality, such as worse water holding capacity, less marbling, less flavor, and reduced storage & processing properties.

The following article will provide detailed information about what meat quality is, how the gut microbiota influences it, and how we can increase meat quality by feeding and modulating the intestinal microflora.

Which factors can contribute to meat quality?

Meat quality is a complex term. On the one hand, meat quality covers measurable parameters such as the content of nutrients, moisture, microbial contamination, etc. On the other hand, and to no small extent, the consumers' preferences are significant. Since meat today is often sold as cuts or in parts (e.g., broiler drumsticks, breast), processing also affects the quality of meat and meat products.

Physical characteristics are objective determinants of meat quality

Physical characteristics are parameters that can be measured. For meat, the following measurable parameters determine meat quality:

1. Fat content and fatty acid composition influence tenderness and taste

Some years ago, the majority of consumers asked for completely lean meat, which, fortunately, has now changed. Fat is a flavor carrier. Especially intramuscular fat (marbling) melts during the preparation, making the meat tender, juicy, and taste good. Fat also transports fat-soluble vitamins.

A further criterion is the composition of the fat, the fatty acids. Geese fat, e.g., is known for its high content of oleic, linoleic, linolenic, and arachidonic acid, all of them derivates of the enzymatic denaturation of stearic acid (<u>Okruszek, 2012</u>).

One exception is cholesterol. Although belonging to the lipids and improving the sensory quality of meat, consumers prefer meat with low cholesterol content.

2. Protein and amino acid content influence the meat value

The content and the composition of protein are important factors in meat quality. Protein is essential for constructing and maintaining organs and muscles and for the functionality of enzymes. The human body needs 20 different amino acids for these tasks, eleven of which it can manufacture by itself. Nine amino acids, however, must be provided by food and are called essential amino acids. Meat is a highly valuable protein source, rich in protein and essential amino acids. The protein quality, therefore, includes the chemical and amino acid score, the index for essential amino acids, and the biological value.

In addition to the pure nutritional value, amino acids contribute to flavor and taste. These flavor amino acids directly influence meat's freshness and flavor and include threonine, alanine, serine, lysine, proline, hydroxyproline, glutamic acid (glutamate is important for the umami taste), aspartic acid, and arginine.

3. Vitamins and trace elements are essential nutrients

Meat is a primary source of B vitamins (B1-B9) and, together with other animal products such as eggs and milk, the only provider of Vitamin B12. Vitamin A is available in the innards, vitamin D in the liver and fat fish, and vitamin K in the flesh.

The most important mineral compounds in meat are zinc, selenium, and iron. Humans can utilize the iron from animal sources particularly well.

4. pH and speed of pH decline decide if the meat is suited for cooking

Since broiler chicken meat nowadays is usually consumed as cut-up pieces or processed products, the appearance at the meat counter or in the plastic box is essential for being sold. The color, seen as an apparent measurement of the freshness and quality of the meat, is influenced by the pH. The muscle pH post-mortem plays an essential role in meat quality. Due to the glycolytic process, the pH post-mortem is a good indication for evaluating physiological meat quality. A rapid pH decline post-mortem to 5.8-6.0 in most cases leads to pale, soft, and exudative (PSE) meat with reduced water retention (Džinić et al., 2015), whereas a high ultimate pH results in dark, firm, and dry (DFD) meat with poor storage quality (Allen et al., 1997)

5. Nobody wants meat like leather

The shear force is a measure of the tenderness of the meat. To determine the shear force, the meat undergoes the process of cooking and chilling. Afterward, standardized meat blocks, with fibers running along the length of the sample, are put into the Warner-Bratzler system. The blade used simulates teeth, and the system measures the force necessary to tear the piece of meat.

6. Microbial contamination is a no-go

The microbial contamination of the meat often occurs during the slaughter process. Let's take a look at salmonella or campylobacter in poultry. The chickens take up salmonella with contaminated feed or water. Campylobacter is transmitted by infected wild birds, inadequately cleaned and disinfected cages, or contaminated water. The bacteria proliferate in the intestine. At slaughter, the intestine's microorganisms can spread onto the meat intended for human consumption.

7. High water holding capacity is necessary to have tender meat

The moisture content contributes to the meat's juiciness and tenderness and improves its quality. If the meat loses its moisture, it gets tough, and quality decreases. Additionally, drip loss reduces the nutritional value of meat and its flavor.

8. Fat oxidation makes meat rancid, and oxidative stress can cause myopathies in broiler breasts

Rancidity of meat occurs when the fat in the flesh gets oxidized. There are different signs of meat rancidity: bad odor, changed color, and a sticky, slimy texture. Poultry meat is considered more susceptible to the development of oxidative rancidity than red meat. This can be explained by its higher content of phospholipids, PUFAs, especially in the thighs. The breast meat, however, has a relatively low level of intramuscular fat (up to 2 %) and, additionally, myoglobin is a natural antioxidant.

But oxidative stress in broiler breasts – and this more and more happens due to a selection of always bigger breasts – can lead to muscle myopathies such as white stripes or wooden breasts, making the meat only usable for processed products.

Sensory meat quality addresses the human senses

Besides physical quality, the sensory and chemical characteristics are essential to meat's economic importance. All attributes of meat that stimulate the human senses (vision, smell, taste, and touch) belong to the sensory quality. It, therefore, is more subjective and hard to determine. The most important features for the consumer include color (attractive or unattractive), texture (tenderness, juiciness, marbling, drip loss), and taste/ flavor (Thorslund et al., 2016).

The appearance is the first impression

Nowadays, meat is often sold as cuts lying in polystyrene or clear plastic trays, over-wrapped with transparent plastic films, so the appearance is paramount. The meat must show an attractive color. Muscle myopathies, such as the ones occurring in chickens, would not meet consumers' needs.

How does the flavor of meat develop?

There is a reaction between reducing sugars and amino acids when meat is cooked (<u>Mottram, 1998</u>). This Maillard reaction, along with the degradation of vitamins, lipid oxidation, and their interaction, is

responsible for the production of the volatile flavor components forming the characteristic aroma and flavor of cooked meat (<u>MacLeod, 1994</u>). <u>Werkhoff et al. (1990</u>) consider cysteine and methionine the most significant contributors to meat flavor development. One factor deteriorating this quality characteristic is lipid peroxidation, which turns the taste to rancid.

Some sensory characteristics are related to physical ones

The parameters of sensory meat quality can be partly explained by measurable parameters. Water retention, e.g., influences the juiciness of the meat. The palatability increases with higher intramuscular fat or marbling (<u>Stewart et al., 2021</u>), the initial pH and the speed of decline decide if the flesh will be pale, soft, and exudative or normal, and lipid peroxidation is the leading cause of a decrease in meat quality (<u>Pereira & Abreu, 2018</u>).

Processing quality

For the processing quality, muscle structure, chemical ingredient interactions, and muscle post-mortem changes are decisive (<u>Berri, 2000</u>).

Does the microbiome influence the meat quality?

The gastrointestinal tract of monogastric animals disposes of a microbiome of primarily bacteria, mainly anaerobic Gram-positive ones (<u>Richards et al., 2005</u>). With its complex microbial community, the digestive tract is responsible for digesting feed and absorbing nutrients, but also for eliminating pathogens and developing immunity. Gut microbiotas play an essential role in digestion, are decisive concerning the synthesis of fatty acids, proteins, and vitamins, and, therefore, influence meat quality (<u>Chen, 2022</u>).

Intestinal microbiotas vary by species/breeds and age (<u>Ma et al., 2022</u>; <u>Sun et al., 2018</u>), and so does meat quality. For example, Duroc pigs with meat of high tenderness, good flavor, and excellent tastiness show different microbiota than other breeds (<u>Xiao, 2017</u>). <u>Zhao et al.(2022</u>) examined high- and low-fat Jinhua pigs, with the high-fat pigs showing more increased backfat thickness but also a higher fat content in the longissimus dorsi. They found low-fat pigs showed a higher abundance of Prevotella and Bacteroides, Ruminococcus sp. AF12-5, Faecalibacterium sp.OFO4-11AC und Oscillibacter sp. CAG:155, which are all involved in fiber fermentation and butyrate production. The high-fat animals showed a higher abundance of Firmicutes and Tenericutes, indicating that they are responsible for higher fat production of the organism in general but also a better fat disposition in the flesh. Lei et al. (2022) showed that abdominal fat was positively correlated with the occurrence of Lachnochlostridium and Christensenelleceae.

The intestinal microbiota-muscle axis enables us to improve meat quality by controlling intestinal microbiota (Lei, 2022). However, to develop strategies to enhance the quality of meat, understanding the composition of the microbiota, the functions of the key bacteria, and the interaction between the host and microbiota is of utmost importance (Chen et al., 2022).

Different factors influence the microbiome

Apart from that microbiotas are different in different breeds, they are additionally influenced by diseases, feeding (diets, medical treatments with, e.g., antibiotics), and the environment (climate, geographical position). This could be shown by different trials. The genetic influence on microbiota was impressively documented by <u>Goodrich et al. (2014)</u>, who detected that the microbiomes of monozygotic twins differ less than the ones of dizygotic twins. Lei et al. (2022) compared the microbiota of two broiler breeds (Arbor Acres and Beijing-You, the last one with a higher abdominal fat rate) and found remarkable differences in

their microbiota composition. When raising them in the same environment and with the same feed, the microbiotas became similar. <u>Zhou et al. (2016)</u> contrasted the cecal microbiota of five Tibetan chickens from five different geographic regions with Lohmann egg-laying hens and Daheng broiler chickens. Besides seeing a difference between the breeds, slightly distinct microbiota between the regions could also be noticed.

The intestinal microbiome can actively be changed by

- promoting the wanted microbes by feeding the appropriate nutrients (e.g., prebiotics)
- reducing the harmful ones by fighting them, for example, with organic acids or phytomolecules
- directly applying probiotics and adding, therefore, desired microbes to the microbiome.

An increase in the abundance of Lactobacillus and Succiniclasticum could be achieved in pigs by feeding them a fermented diet, and Mitsuokella and Erysipelotrichaceae proliferated by adding a probiotic containing B. subtilis and E. faecalis to the diet (<u>Wang et al., 2022</u>).

How to change the intestinal microbiome to improve meat quality?

Before changing the microbiome, we must know which microbes are "responsible" for which characteristics. However, the microbiotas do not act individually but as consortia. The following table shows a selection of bacteria that, besides supporting the gut and its functions, influence meat quality in some way.

Metabolites	Producing bacteria	Biological functions and effects on pigs
Short-chain fatty acids (acetate, butyrate, and propionate)	Ruminococcaceae Ruminococcus Lachnospiraceae Blautia Roseburia Lactobacillaceae Clostridium Eubacterium Faecalibacterium Bifidobacterium Bacteroides	Regulate lipid metabolism Improve meat quality
Lactate	Lactic acid bacteria Bifidobacterium	Important metabolite for cross-feeding of SCFA-producing microbiota
Bile acids (primary and secondary bile acids)	Clostridium species Eubacterium Parabacteroides Lachnospiraceae	Regulate lipid metabolism
Ammonia	Amino acid fermenting commensals Helicobacter	By-product of amino acid fermentation Inhibits short-chain fatty acid oxidation
B Vitamins and vitamin K	Bacteroides Lactobacillus	Serve as coenzymes in neurological processes (B vitamins) • Essential vitamin for proper blood clotting (vitamin K)

Table 1: Bacteria influencing meat quality (according to Vasquez et al., 2022)

Fat for meat quality is intramuscular fat

If we talk about increasing fat to improve meat quality, we talk about increasing intramuscular fat or marbling, not depot fat. The fat in meat-producing animals is mostly a combination of triglycerides from the diet and fatty acids synthesized. Fat deposition and composition in non-ruminants reflect the fatty acid composition of the diet but are also closely related to the design of the microbiome; short-chain fatty acids in monogastric, e.g., are exclusively produced by the gut microbiome (Dinh et al., 2021; Vasquez et al., 2022). Intramuscular fat is mainly made of triglycerides but also disposes of phospholipids associated with proteins, such as lipoproteins or proteolipids, influencing meat flavor. The fermentation of indigestible polysaccharides or amino acids results in short-chain or branched-chain fatty acids, respectively. Lactate, produced by lactic acid bacteria, is utilized by SCFA-producing microbiota. An imbalance in the microbiome fosters lipid deposition, as shown by Kallus and Brandt (2012), who found a higher proportion of Firmicutes to Bacteroidetes (50% higher) in obese mice than in lean ones. In a trial described by Zhou et al. (2016), tiny Tibetian chickens with a low percentage of abdominal fat were compared to two breeds (Lohmann layers and Daheng broilers) being large and with a high percentage of abdominal fat. The Tibetan chickens showed a two to four-fold higher abundance of Christensenellacea in the cecal microbiome. Christensenellas belong to the bacterial strain of firmicutes. They are linked to slimness in human nutrition, which was already proven by Goodrich et al. (2014) and is the contrary stated by Lei et al. (2022).

Another example was provided by <u>Wen et al. (2023)</u>. They compared two broiler enterotypes distinguished by Clostridia vadinB60 and Rikenellaceae_RC9_gut and saw that the type with an abundance of Clostridia_vadinBB60 showed higher intramuscular fat content but also more subcutaneous fat tissue. The scientists also found another bacterium especially responsible for intramuscular fat: A lower plethora of Clostridia vadimBE97 resulted in a higher intramuscular fat content in breast and thigh muscles but not adipose tissues. Similar results were achieved in a trial with pigs and mice: Jinhua pigs showed a significantly higher level of intramuscular fat than Landrace pigs. When transplanting the fecal microbiota of the two breeds in mice, the mice showed similar characteristics in fat metabolism as their donors of feces (<u>Wu et al., 2021</u>).

According to several studies (e.g., <u>Chen et al., 2008</u>; <u>Liu et al., 2019</u>), intramuscular fat in chicken has a low heritability but may be controlled by feeding up to a certain extent. In pigs, <u>Lo et al. (1992</u>) and <u>Ding et al. (2019</u>) found a moderate to low (0.16 – 0.23) heritability for intramuscular fat, but Cabling et al. (2015) calculated a heritability of 0.79 for the marbling score.

At least, especially the composition of fatty acids can easily be changed in monogastric (<u>Aaslyng and</u> <u>Meinert, 2017</u>). <u>Zou et al. (2017</u>) examined the effect of Lactobacillus brevis and tea polyphenol, each alone or combining both. Lactobacillus is probably involved in turning complex carbohydrates into metabolites lactose and ethanol, but also acetic acid and SCFA. SCFAs are mainly produced by Saccharolytic and anaerobic microbiota, aiding in the degradation of carbohydrates the host cannot digest (e.g., cellulose or resistant polysaccharides into monomeric and dimeric sugars and fermenting them subsequently into short-chain fatty acids). Including fibers and various oligosaccharides was shown to increase the gut microbiome's fermentation capacity for producing short-chain fatty acids.

In a trial conducted by Jiao et al. (2020), they showed that SCFAs applied in the ileum modulate lipid metabolism and lead to higher meat quality in growing pigs. A plant polyphenol was used by <u>Yu et al.</u> (2021). The added resveratrol, a plant polyphenol in grapes and grape products, to the diet of Peking ducks and could significantly increase intramuscular fat.

Oxidation of lipids and proteins must be prevented

The composition of the fatty acids and occurring oxidative stress in adipose and muscle tissue influences or impacts meat quality in farm animals (<u>Chen et al., 2022</u>). During the last few years, the demand for healthier animal products containing higher levels of polyunsaturated fatty acids has increased. Consequently, the risk of lipoperoxidation has risen (<u>Serra et al., 2021</u>). Solutions are needed to counteract this deterioration of meat quality. As can be seen in table 1, ammonia produced by amino acid-fermenting commensals and Helicobacter inhibits the oxidation of SCFAs. <u>Ma et al. (2022</u>) changed the microbiome of sows by feeding a probiotic from mating till day 21 of lactation and achieved a decreased level of MDA, a

sign of reduced oxidative stress. Similar results were achieved by <u>He et al. (2022)</u>. In their trial, the supplementation of 200 mg yeast ß-glucan/kg of feed significantly decreased the abundance of the phylum WPS-2 as well as markedly increased catalase, superoxide dismutase (both p<0.05) and the total antioxidant activity (p<0.01) in skeletal muscle. Another approach was done by <u>Wu et al. (2020)</u> in broilers. They applied glucose oxidases (GOD) produced by Aspergillus niger and Penicillium amagasakiense. Both enzymes did not disturb but improved beneficial bacteria and microbiota. The GOD produced by A. niger reduced the content of malondialdehyde in the plasma.

Another alternative is antioxidant extracts from plants (<u>Džinić, 2015</u>). As consumers nowadays bet more on natural products, they would be good candidates. They are considered safe and, therefore, wellaccepted by consumers and have beneficial effects on animal health, welfare, and production performance.

<u>Hazrati et al. (2020)</u> showed in a trial that the essential oils of ajwain and dill decreased the concentration of malondialdehyde (MDA) in quails' breast meat and, therefore, lipid peroxidation and reduced cooking loss. The antioxidant effects of thymol and carvacrol were shown by <u>Luna et al. (2010)</u>. The group receiving the essential oils showed lower TBARS in the thigh samples than the control group but similar TBARS to the butylated hydroxytoluene-provided group.

Protein quality is a question of essential amino acids

Protein with a high content of essential amino acids is one of the most critical components of meat. Alfaig et al. (2014) tested probiotics and thyme essential oil in broilers. They found out that the content of EAAs in breast and thigh muscles numerically increased gradually from the control over the probiotic and a combination of a probiotic up to the thyme essential oil group. A significant (p<0.05) increase in all tested amino acids (arginine, cysteine, phenylalanine, histidine, isoleucine, leucine, lysine, methionine, threonine, and valine) could be observed in the samples of the breast and the thigh muscles when comparing the thyme essential oil group with the control. Zou et al. (2017) provided similar results, showing a significant increase in leucine and glutamic acid as well as a numerical increase in lysin, valine, methionine, isoleucine, phenylalanine, threonine, asparagine, alanine, glycin, serin, and proline through the addition of a combination of Lactobacillus brevis and tea polyphenols. They also determined an increase in the beneficial bacteria Lactobacillus and Bacteroides. The experimental results led them to the assumption that both additives may also improve the taste of meat by increasing some of the essential and delicate flavors produced by amino acids.

Tenderness is closely related to drip loss

The already mentioned trial conducted by Lei et al. (2022) with two different broiler breeds (Arbor Acres and Beijing-You) having different microbiota showed a negative correlation between drip loss and the abundance of Lachnochlostridium. They remodeled the Arbor Acres' microbiome by applying a bacterial suspension derived from the Beijing-You breed and decreased drip loss in their meat. <u>He et al. (2022)</u> changed the microbiome by adding yeast ß-glucan to the diet of finisher pigs. They achieved a reduced cooking loss (linear, p<0.05) and a lower drip loss (p<0.05), together indicating a better water-holding capacity, as well as a decreased lactate content. The addition of a multi-species probiotic to the diet of finishing pigs tended to result in lower cooking and drip loss(p<0.1) besides modulating the intestinal flora (higher lactobacilli and lower E. coli counts in the feces) (Balasubramanian et al., 2017) and the inclusion of Lactobacillus brevis and tea polyphenol individually or in a synergistic combination improved water holding capacity and decreased drip loss Zou et al. (2017).

<u>Puvača et al. (2019</u>) observed the lowest drip-loss values in breast meat and thigh with drumstick through feeding chickens 0.5 g or 1.0 g of hot red pepper per 100 g of feed, respectively, in the grower and finisher phase. The feeding of resveratrol reduced drip loss of Peking ducks' leg muscles. SCFA infused into the ileum enlarged the longissimus dorsi area and alleviated drip loss (Jiao et al, 2021).

The decrease and increase of the pH after slaughtering determines meat quality

The pH in the muscles of a living animal is about 7.2. With slaughtering and bleeding, the energy supply of the muscles is interrupted. The stored glycogen gets degraded to lactic acid, lowering the pH. Usually, the lowest pH value of 5.4-5.7 in meat is reached after 18 to 24 hours. Afterward, it starts to rise again.

In stressed animals, the stress hormones adrenalin and noradrenalin provoke a rushly occurring and, due to a lack of oxygen, anaerobic metabolism and the quick production of lactic acid. This too rapid decrease in pH leads to the denaturation of proteins in the muscle cells and reduced water-holding capacity. The result is PSE (pale, soft, and exudative) meat.

On the contrary, DFD meat (dark, firm, and dry) occurs if the glycogen reserves, due to challenges, are already used up, and the lactic acid production is insufficient. Especially PSE meat is closely related to breeds – some are more susceptible to stress, others less. However, some trials show that influencing pH in meat is possible to a certain extent.

<u>He et al., 2022</u> added yeast ß-glucan to the diets of finishing pigs and a higher $pH_{45 \text{ min}}$ (linear and quadratic, p<0.01) and a higher redness (a*; linear, p<0.05) of the meat. <u>Wu et al. (2020)</u> achieved a significantly increased pH_{24h} through the addition of Glucose oxidase produced by Aspergillus niger.

Sensory characteristics are very subjective

In general, the sensory characteristics of meat are seen very individually. Some prefer lean, others fatty meat, some like meat with a characteristic taste, and others with a neutral. However, the typical meat taste of umami is partly determined by the nucleotide inosine monophosphate (IMP), which is regarded as an essential index for evaluating meat flavor and the acceptability of meat products. IMP provides about 40-fold higher umami taste than sodium glutamate (Huang et al. 2022). IMP is the organophosphate of inosin. Inosine, however, according to Kroemer and Zitvogel (2020), is produced by Bifidobacterium pseudolongum, which possibly can be controlled by feeding. Sun et al. (2018) compared Caoke and Partridge Shank chickens and divided them into free-range and cage groups. They found out that, except for acids, the amounts of flavor components were higher in the free-range than in the cage groups. The two housing systems also modified the microbiota, and Sun et al. took it as an indication that meat flavor, as well as the composition and diversity of gut microbiota, are closely associated with the housing systems. Fu et al. (2023) examined the addition of a mixture containing Pulsatilla, Gentian, and Rhizoma coptidis and a mixture with Codonopsis pilosula, Atractylodes, Poria cocos, and Licorice to the feed of Hungarian white geese. They saw that in both groups, the total amino acid levels, especially Glu, Lys, and Asp, increased, with, according to Liu et al. (2018), Glu and Asp directly affecting meat's freshness and flavor. Yu et al. (2021) achieved similar results by adding resveratrol to the diet of Peking ducks. The addition of the herbs additionally led to a higher Firmicutes/Bacteroidetes ratio and an increased level of lactobacilli (Fu et al., 2023).

How can EW Nutrition's feed additives help to improve meat quality?

Meat quality is influenced by the microbiome. So, feed additives that stabilize the microbiome or promote certain beneficial bacterial strains are an opportunity.

Ventar D modulates the microbiome

Ventar D balances the microbiome by promoting beneficial bacteria such as lactobacilli and fighting harmful ones such as Clostridia, E. coli, and Salmonella. (<u>Heinzl, 2022</u>). In another trial with broilers, the addition of Ventar D to all feeds (100 g/t) showed an increase in short-chain fatty acids in the intestine:



Figure 1: Short-chain fatty acids in the cecum of broilers

Santoquin countersteers oxidation

Another helpful product category is antioxidants. They can prevent the oxidation of lipids and proteins. For this purpose, EW Nutrition offers Santoquin M6*, a product tested by Kuttapan et al. (2021). Santoquin M6 was tested concerning its ability to minimize the oxidative damage caused by feeding oxidized fat. A control group receiving oxidized fat in feed was compared to one receiving oxidized fat plus 188 ppm Santoquin M6 (\triangleq 125 ppm ethoxyquin). The main parameters for this study were TBARS in the breast muscle, the incidence of wooden breast, and the live weight on day 48.

Results indicated that the inclusion of Santoquin M6 reduced the production of TBARS in the breast muscles, demonstrating a lower level of oxidative stress in the breast muscles.



Figure 2: Thiobarbituric acid reactive substances (TBARS) in broiler breast muscles. TBARS are formed as a byproduct of lipid peroxidation.

Additionally, it reduced the incidence of severe woody breasts (Score 3) by almost half and helped

mitigate the impact of breast muscle degradation due to increased oxidative stress.



Figure 3: Incidence of wooden breast in broilers

*Usage of ethoxyquin is dependent on country regulations.

Feed hygiene with Acidomix products minimizes harmful pathogens

The Acidomix product line offers liquid, powdery, and micro-granulated products to be added to feed and water. The organic acids in Acidomix directly act against pathogens in the feed and the water and help keep the intestinal flora in balance.

A trial evaluating the effect of different Acidomix products against diverse pathogens showed lower MICs for most Acidomix products than for single organic acids. The trial was conducted with decreasing

concentrations of the Acidomix products (2 – 0.015625 %) and 10^5 CFU of the respective microorganisms (microtiter plates; 50 µl bacterial solution and 50 µl diluted product).

Feeding is the one side, slaughtering the other one

With feeding, the microbiota and some meat characteristics can be changed; however, the last step, handling the animals before and the meat after slaughtering also significantly contributes to a good quality of meat. Stress due to the transport and the slaughterhouse atmosphere, combined with stress-sensible breeds, can lead to PSE meat. Incorrect handling at the slaughterhouse can lead to meat contaminated with pathogens.

Combining feeding measures with professional and calm handling of the animals is the best strategy to achieve high-quality meat.

References

0.6

Aaslyng, Margit D., and Lene Meinert. "Meat Flavour in Pork and Beef – from Animal to Meal." *Meat Science* 132 (2017): 112–17. https://doi/org/10.1016/jnscatsgit2019.04.0122ainst different pathogens (%)

Alfaig, Ebrahim, Maria Angelovičova, Martin Kral, and Ondrej Bučko. "EFF ECT of Probiotics and Thyme Essential Oil on the Essential Amino Acid Content of the Broiler Chicken Meat." Acta Scientiarum Polonorum Technologia Alimentaria 13, no.º4 (2014): 425–32. https://doi.org/10.17306/j.afs.2014.4.9.

Allen, CD, SM Russell, and DL Fletcher. "The Relationship of Broiler Breast Meat Color and Ph to Shelf-Life and Odor Development?" *Poultry Science* 76, no. 7 (1997): 1042–46. https://doi.org/10.1093/ps/76.7.1042.

^{0.1} Balasubramanian, Balamuralikrishnan, Sang In Lee, and In-Ho Kim. "Inclusion of Dietary Multi-Species Probiotic on Growth Performance, Nutrient Digestibility, Meat Quality Traits, Faecal Microbiota, and Diarrhoea Score in Growing-Finishing Pigs." Italian Journal of Animal Science 17, no. 1 (2017): 100–106. https://doi.org/10.1080/1828051X.2017.1340097 Acidomix FL

Berri, Céile. "Variability of Sensory and Processing Qualities of Poultry Meat." World's Poultry Science Journal 56, no. 3 (2000): 209–24. https://doi.org/10.1079/wps20000016.

Cabling, M. M., H. S. Kang, B. M. Lopez, M. Jang, H. S. Kim, K. C. Nam, J. G. Choi, and K. S. Seo. "Estimation of Genetic Associations between Production and Meat Quality Traits in Duroc Pigs." Asian-Australasian Journal of Animal Sciences 28, no. 8 (2015): 1061–65. https://doi.org/10.5713/ajas.14.0783.

Chen, Binlong, Diyan Li, Dong Leng, Hua Kui, Xue Bai, and Tao Wang. "Gut Microbiota and Meat Quality." *Frontiers in Microbiology* 13 (2022). https://doi.org/10.3389/fmicb.2022.951726.

Chen, J.L., G.P. Zhao, M.Q. Zheng, J. Wen, and N. Yang. "Estimation of Genetic Parameters for Contents of Intramuscular Fat and Inosine-5´-Monophosphate and Carcass Traits in Chinese Beijing-You Chickens." *Poultry Science* 87, no. 6 (2008): 1098–1104. <u>https://doi.org/10.3382/ps.2007-00504</u>.

Ding, Rongrong, Ming Yang, Jianping Quan, Shaoyun Li, Zhanwei Zhuang, Shenping Zhou, Enqin Zheng, et al. "Single-Locus and Multi-Locus Genome-Wide Association Studies for Intramuscular Fat in Duroc Pigs." Frontiers in Genetics 10 (2019). https://doi.org/10.3389/fgene.2019.00619.

Dinh, Thu T., K. Virellia To, and M. Wes Schilling. "Fatty Acid Composition of Meat Animals as Flavor Precursors." *Meat and Muscle Biology* 5, no. 1 (2021). https://doi.org/10.22175/mmb.12251.

Džinić, N., N. Puvača, T. Tasić, P. Ikonić, and Okanović. "How Meat Quality and Sensory Perception Is Influenced by Feeding Poultry Plant Extracts." *World's Poultry Science Journal* 71, no. 4 (2015): 673–82. https://doi.org/10.1017/s0043933915002378.

Džinić, N., N. Puvača, T. Tasić, P. Ikonić, and Okanović. "How Meat Quality and Sensory Perception Is Influenced by Feeding Poultry Plant Extracts." *World's Poultry Science Journal* 71, no. 4 (2015): 673–82. https://doi.org/10.1017/s0043933915002378.

Fu, Guilin, Yuxuan Zhou, Yupu Song, Chang Liu, Manjie Hu, Qiuyu Xie, Jingbo Wang, et al. "The Effect of Combined Dietary Supplementation of Herbal Additives on Carcass Traits, Meat Quality, Immunity and Cecal Microbiota Composition in Hungarian White Geese (v0.2)"." *Peer J.;* 11:e15316, May 8, 2023. https://doi.org/10.7287/peerj.15316v0.2/reviews/3.

Fu, Guilin, Yuxuan Zhou, Yupu Song, Chang Liu, Manjie Hu, Qiuyu Xie, Jingbo Wang, et al. "The Effect of Combined Dietary Supplementation of Herbal Additives on Carcass Traits, Meat Quality, Immunity and Cecal Microbiota Composition in Hungarian White Geese." *PeerJ* 11 (2023). https://doi.org/10.7717/peerj.15316.

Goodrich, Julia K., Jillian L. Waters, Angela C. Poole, Jessica L. Sutter, Omry Koren, Ran Blekhman, Michelle Beaumont, et al. "Human Genetics Shape the Gut Microbiome." *Cell* 159, no. 4 (2014): 789–99. https://doi.org/10.1016/j.cell.2014.09.053.

Hazrati, S., V. Rezaeipour, and S. Asadzadeh. "Effects of Phytogenic Feed Additives, Probiotic and Mannan-Oligosaccharides on Performance, Blood Metabolites, Meat Quality, Intestinal Morphology, and Microbial Population of Japanese Quail." *British Poultry Science* 61, no. 2 (2019): 132–39. https://doi.org/10.1080/00071668.2019.1686122.

He, Linjuan, Jianxin Guo, Yubo Wang, Lu Wang, Doudou Xu, Enfa Yan, Xin Zhang, and Jingdong Yin. "Effects of Dietary Yeast β -Glucan Supplementation on Meat Quality, Antioxidant Capacity and Gut Microbiota of Finishing Pigs." Antioxidants 11, no. 7 (2022): 1340. https://doi.org/10.3390/antiox11071340.

Heinzl, Inge. "Efficient Microbiome Modulation with Phytomolecules." EW Nutrition, July 6, 2022. https://ew-nutrition.com/pushing-microbiome-in-right-direction-phytomolecules/.

Huang, Zengwen, Juan Zhang, Yaling Gu, Zhengyun Cai, Dawei Wei, Xiaofang Feng, and Chaoyun Yang. "Analysis of the Molecular Mechanism of Inosine Monophosphate Deposition in Jingyuan Chicken Muscles Using a Proteomic Approach." *Poultry Science* 101, no. 4 (2022): 101741. https://doi.org/10.1016/j.psj.2022.101741.

Jiao, Anran, Hui Diao, Bing Yu, Jun He, Jie Yu, Ping Zheng, Yuheng Luo, et al. "Infusion of Short Chain Fatty Acids in the Ileum Improves the Carcass Traits, Meat Quality and Lipid Metabolism of Growing Pigs." *Animal Nutrition* 7, no. 1 (2021): 94–100. https://doi.org/10.1016/j.aninu.2020.05.009.

Kallus, Samuel J., and Lawrence J. Brandt. "The Intestinal Microbiota and Obesity." *Journal of Clinical Gastroenterology* 46, no. 1 (2012): 16–24. https://doi.org/10.1097/mcg.0b013e31823711fd.

Khan, Muhammad Issa, Cheorun Jo, and Muhammad Rizwan Tariq. "Meat Flavor Precursors and Factors Influencing Flavor Precursors—a Systematic Review." *Meat Science* 110 (2015): 278–84. https://doi.org/10.1016/j.meatsci.2015.08.002.

Kroemer, Guido, and Laurence Zitvogel. "Inosine: Novel Microbiota-Derived Immunostimulatory Metabolite." *Cell Research* 30, no. 11 (2020): 942–43. https://doi.org/10.1038/s41422-020-00417-1.

Kuttappan, Vivek A., Megharaja Manangi, Matthew Bekker, Juxing Chen, and Mercedes Vazquez-Anon. "Nutritional Intervention Strategies Using Dietary Antioxidants and Organic Trace Minerals to Reduce the Incidence of Wooden Breast and Other Carcass Quality Defects in Broiler Birds." Frontiers in Physiology 12 (2021). https://doi.org/10.3389/fphys.2021.663409

Lei, Jiaqi, Yuanyang Dong, Qihang Hou, Yang He, Yujiao Lai, Chaoyong Liao, Yoichiro Kawamura, Junyou Li, and

Bingkun Zhang. "Intestinal Microbiota Regulate Certain Meat Quality Parameters in Chicken." *Frontiers in Nutrition* 9 (2022). https://doi.org/10.3389/fnut.2022.747705.

Liu, R., M. Zheng, J. Wang, H. Cui, Q. Li, J. Liu, G. Zhao, and J. Wen. "Effects of Genomic Selection for Intramuscular Fat Content in Breast Muscle in Chinese Local Chickens." *Animal Genetics* 50, no. 1 (2018): 87–91. https://doi.org/10.1111/age.12744.

Lo, L. L., D. G. McLaren, F. K. McKeith, R. L. Fernando, and J. Novakofski. "Genetic Analyses of Growth, Real-Time Ultrasound, Carcass, and Pork Quality Traits in Duroc and Landrace Pigs: II. Heritabilities and Correlations." Journal of Animal Science 70, no. 8 (1992): 2387–96. https://doi.org/10.2527/1992.7082387x.

Luna, A., M.C. Lábaque, J.A. Zygadlo, and R.H. Marin. "Effects of Thymol and Carvacrol Feed Supplementation on Lipid Oxidation in Broiler Meat." *Poultry Science* 89, no. 2 (2010): 366–70. https://doi.org/10.3382/ps.2009-00130.

Ma, Cui, Md. Abul Azad, Wu Tang, Qian Zhu, Wei Wang, Qiankun Gao, and Xiangfeng Kong. "Maternal Probiotics Supplementation Improves Immune and Antioxidant Function in Suckling Piglets via Modifying Gut Microbiota." *Journal of Applied Microbiology* 133, no. 2 (2022): 515–28. https://doi.org/10.1111/jam.15572.

Ma, Jianfeng, Jingyun Chen, Mailin Gan, Lei Chen, Ye Zhao, Yan Zhu, Lili Niu, Shunhua Zhang, Li Zhu, and Linyuan Shen. "Gut Microbiota Composition and Diversity in Different Commercial Swine Breeds in Early and Finishing Growth Stages." *Animals* 12, no. 13 (2022): 1607. https://doi.org/10.3390/ani12131607.

MacLeod, G. "The Flavour of Beef." Essay. In *Shahidi, F. (Eds) Flavor of Meat and Meat Products*, 4–37. Boston, MA: Springer, 1994.

Mottram, Donald. "Flavour Formation in Meat and Meat Products: A Review." Food Chemistry 62, no. 4 (1998): 415-24. https://doi.org/10.1016/s0308-8146(98)00076-4.

Okruszek, A. "Fatty Acid Composition of Muscle and Adipose Tissue of Indigenous Polish Geese Breeds." *Archives Animal Breeding* 55, no. 3 (2012): 294–302. https://doi.org/10.5194/aab-55-294-2012.

Pereira, Ana Lúcia F., and Virgínia Kelly G. Abreu. "Lipid Peroxidation in Meat and Meat Products." Essay. In *Lipid Peroxidation Research*. London: IntechOpen, 2020.

Puvača, Nikola, Tatjana Peulić, Predrag Ikonić, Sanja Popović, Jasmina Lazarević, Olivera Đuragić, Magdalena Cara, and Nedeljka Nikolova. "Effects of Medicinal Plants in Broiler Chicken Nutrition on Selected Parameters of Meat Quality." *Macedonian Journal of Animal Science* 9, no. 2 (2019): 45–51. https://doi.org/10.54865/mjas1992045p.

Richards, J. D., J. Gong, and C. F. de Lange. "The Gastrointestinal Microbiota and Its Role in Monogastric Nutrition and Health with an Emphasis on Pigs: Current Understanding, Possible Modulations, and New Technologies for Ecological Studies." Canadian Journal of Animal Science 85, no. 4 (2005): 421–35. https://doi.org/10.4141/a05-049.

Serra, Valentina, Giancarlo Salvatori, and Grazia Pastorelli. "Dietary Polyphenol Supplementation in Food Producing Animals: Effects on the Quality of Derived Products." *Animals* 11, no. 2 (2021): 401. https://doi.org/10.3390/ani11020401.

Stewart, S.M., G.E. Gardner, P. McGilchrist, D.W. Pethick, R. Polkinghorne, J.M. Thompson, and G. Tarr. "Prediction of Consumer Palatability in Beef Using Visual Marbling Scores and Chemical Intramuscular Fat Percentage." *Meat Science* 181 (2021): 108322. https://doi.org/10.1016/j.meatsci.2020.108322.

Sun, Jing, Yan Wang, Nianzhen Li, Hang Zhong, Hengyong Xu, Qing Zhu, and Yiping Liu. "Comparative Analysis of the Gut Microbial Composition and Meat Flavor of Two Chicken Breeds in Different Rearing Patterns." *BioMed Research International* 2018 (2018): 1–13. https://doi.org/10.1155/2018/4343196.

Thorslund, Cecilie A.H., Peter Sandøe, Margit Dall Aaslyng, and Jesper Lassen. "A Good Taste in the Meat, a Good Taste in the Mouth – Animal Welfare as an Aspect of Pork Quality in Three European Countries." *Livestock Science* 193 (2016): 58–65. https://doi.org/10.1016/j.livsci.2016.09.007.

Vasquez, Robie, Ju Kyoung Oh, Ji Hoon Song, and Dae-Kyung Kang. "Gut Microbiome-Produced Metabolites in Pigs: A Review on Their Biological Functions and the Influence of Probiotics." *Journal of Animal Science and Technology* 64, no. 4 (2022): 671–95. https://doi.org/10.5187/jast.2022.e58.

Wang, Cheng, Siyu Wei, Bojing Liu, Fengqin Wang, Zeqing Lu, Mingliang Jin, and Yizhen Wang. "Maternal Consumption of a Fermented Diet Protects Offspring against Intestinal Inflammation by Regulating the Gut Microbiota." *Gut Microbes* 14, no. 1 (2022). https://doi.org/10.1080/19490976.2022.2057779.

Wen, Chaoliang, Qinli Gou, Shuang Gu, Qiang Huang, Congjiao Sun, Jiangxia Zheng, and Ning Yang. "The Cecal Ecosystem Is a Great Contributor to Intramuscular Fat Deposition in Broilers." *Poultry Science* 102, no. 4 (2023): 102568. https://doi.org/10.1016/j.psj.2023.102568.

Werkhoff, Peter, Juergen Bruening, Roland Emberger, Matthias Guentert, Manfred Koepsel, Walter Kuhn, and Horst Surburg. "Isolation and Characterization of Volatile Sulfur-Containing Meat Flavor Components in Model Systems." *Journal of Agricultural and Food Chemistry* 38, no. 3 (1990): 777–91. https://doi.org/10.1021/jf00093a041.

Wu, Choufei, Wentao Lyu, Qihua Hong, Xiaojun Zhang, Hua Yang, and Yingping Xiao. "Gut Microbiota Influence Lipid Metabolism of Skeletal Muscle in Pigs." *Frontiers in Nutrition* 8 (2021). https://doi.org/10.3389/fnut.2021.675445.

Xiao, Yingping, Kaifeng Li, Yun Xiang, Weidong Zhou, Guohong Gui, and Hua Yang. "The Fecal Microbiota Composition of Boar Duroc, Yorkshire, Landrace and Hampshire Pigs." *Asian-Australasian Journal of Animal Sciences* 30, no. 10 (2017): 1456–63. https://doi.org/10.5713/ajas.16.0746.

Yu, Qifang, Chengkun Fang, Yujing Ma, Shaoping He, Kolapo Matthew Ajuwon, and Jianhua He. "Dietary Resveratrol Supplement Improves Carcass Traits and Meat Quality of Pekin Ducks." *Poultry Science* 100, no. 3 (2021): 100802. https://doi.org/10.1016/j.psj.2020.10.056.

Zhao, Guangmin, Yun Xiang, Xiaoli Wang, Bing Dai, Xiaojun Zhang, Lingyan Ma, Hua Yang, and Wentao Lyu. "Exploring the Possible Link between the Gut Microbiome and Fat Deposition in Pigs." *Oxidative Medicine and Cellular Longevity* 2022 (2022): 1–13. https://doi.org/10.1155/2022/1098892.

Zhou, Xueyan, Xiaosong Jiang, Chaowu Yang, Bingcun Ma, Changwei Lei, Changwen Xu, Anyun Zhang, et al. "Cecal Microbiota of Tibetan Chickens from Five Geographic Regions Were Determined by 16s Rrna Sequencing." *MicrobiologyOpen* 5, no. 5 (2016): 753-62. https://doi.org/10.1002/mbo3.367.

Zou, Xiaozhuo, Rong Xiao, Huali Li, Ting Liu, Yong Liao, Yuanliang Wang, Shusong Wu, and Zongjun Li. "Effect of a Novel Strain of *Lactobacillus Brevis* M8 and Tea Polyphenol Diets on Performance, Meat Quality and Intestinal Microbiota in Broilers." *Italian Journal of Animal Science* 17, no. 2 (2017): 396–407. https://doi.org/10.1080/1828051x.2017.1365260.

DDGS and oxidative damage



Distiller's dried grains are produced by condensing and drying the stillage left over after starchfermentation of corn for ethanol production. Solubles left over from the process are usually added before drying, resulting in the DDGS product that has become more and more commonplace in poultry feed formulations.

Historically, this ingredient was used in ruminant diets, as the nutrient content and quality of DDGS is considered to be somewhat variable – not only between, but also within production facilities. However, recent improvements in processing technologies and quality control systems have resulted in more consistent, higher-quality feed products.

Furthermore, newer fractionation processes continue to be implemented by some ethanol plants that are capable of fractioning out protein, fiber and oil portions of the grain, either pre- or post-fermentation, providing a wide variety of co-products which will result from the blending of these fraction streams.

Why is there more danger of oxidative damage today?

A decade ago, typical poultry diets consisted of far fewer feed ingredients than some of today's rations. Additionally, the use of further-processed by-product meals and fats has also increased due to economic constraints associated with some raw ingredients.

This has created a diet which is highly prone to oxidative degeneration due to the higher rate of thermal, and in some cases, chemical processing of feed ingredients such as DDGS.

Currently, it is not uncommon to see DDGS inclusion rates between 5 and 12% in broiler and turkey diets, depending on bird age and feed prices.

One factor to consider when utilizing DDGS at such levels is the increase in polyunsaturated fatty acid content relative to saturated fatty acids that results from the addition of the 18:2, n-6 -rich corn oil which comprises approximately 10% of most DDGS.

Despite the fact that most modern rations have decreased the total lipid content in recent years, overall increases in the use of vegetable-based by-products invariably change the fatty acid profile of the diet to one which contains higher levels of polyunsaturated fats relative to saturated fats.

Polyunsaturated fatty acids are highly sensitive to oxidation during storage and are likely to turn rancid at high environmental temperatures. The oxidation process involves the generation of fatty acid free radicals, which then react with molecular oxygen to produce peroxide free radicals. This results in a chemical change to the fatty acid which decreases nutrient value and often produces undesirable odor.

...And that's why risk management of oxidative damage is essential

Oxidative damage in feeds entails economic losses because of the negative impact on feed quality through the transformation of the lipid fraction of feed ingredients, decreased animal growth rate and performance, and decreased meat quality parameters, such as nutritional value and shelf-life. Therefore, lipid stability in feed is important, particularly with regard to the oxidative rancidity that occurs in high-fat ingredients, and prevention and management of oxidative stress is critical.

Respiratory health in poultry: no action is no solution



by Inge Heinzl and Ruturaj Patil, EW Nutrition

Broilers face high respiratory disease risks. In winter, they often come from lower temperatures; throughout the year, they come from improper ventilation and proximity to manure or infected birds. The confined spaces and lack of proper airflow create an environment conducive to harmful airborne particles

and pathogens, significantly compromising birds' respiratory health. In the possible presence of viruses such as ILT (Infectious Laryngotracheitis Virus), IBV (Infectious Bronchitis Virus), AIV (Avian Influenza Virus, NDV (Newcastle Disease Virus), bacteria like *Mycoplasma gallisepticum*, *E. coli*, or Chlamydia, respiratory issues are inevitable.

High efficiency takes its toll

A bird, generally a flying species, has a complex respiratory system. Instead of the diaphragm cooperating with the lung, nine additional air sacs do the job of sucking in and blowing out of the air like bellows. They increase the air volume passing through the lungs, where oxygen absorption occurs. The air sacs are situated in different parts of the birds and connected to hollow (pneumatic) bones.

The co-action of the air sacs and the lung results in a high efficiency of the bird's respiratory system: birds can extract about 160% more oxygen from the air than mammals. However, the extended parts of the respiratory system also offer a high contact surface for pathogens. To protect themselves, the respiratory system is equipped with

- cilia in the trachea to propel entrapped particles for disposal

- mucus produced by goblet cells in the trachea and cooperating with the cilia
- immune cells in the lung, scavenging inhaled particles and bacteria that enter the lower respiratory tract

Additional support is recommended

To additionally support your birds against respiratory issues, stress should be kept low, and immunity to diseases should be high. If possible, decrease the stocking density. Effective litter management can help keep litter particle inhalation low. These particles irritate the respiratory system and reduce immune resistance. They often carry pathogens and possibly induce respiratory issues through several toxic mechanisms.

Another possibility is using phytogenic substances alone or combined with vaccines. Eucalyptus oil exerts antimicrobial, anti-inflammatory, mucolytic, and bronchodilator effects in the case of respiratory disease. Thyme has expectorant, mucolytic, antitussive, and antispasmodic characteristics, and mint, with its antihistamine and cooling effect, acts as a decongestant. <u>Grippozon</u> is such an example, based on fast-acting, concentrated phytomolecules supporting animals against respiratory challenges.



Gurgling sounds (post-vaccine, % cages)

A trial with 20,000 birds showed fewer gurgling sounds and reduced post-vaccination reaction than the untreated group.

Regardless of the solution chosen, especially with the cold season coming and high stocking density a

Feeding layers for longer laying cycles and optimized production



Conference report

At the recent EW Nutrition Poultry Academy in Jakarta Indonesia, Dr Steve Leeson, Professor Emeritus, University of Guelph, Canada, commented that "genetic progress in layer breeding has been substantial in recent decades. Since 1995, the <u>yearly</u> change has included +1 egg, -0.01 feed/dozen eggs, -10g final bodyweight, 0.02% mortality, and +1 week at >90% egg production. This improved persistency of commercial laying hens enables egg producers to keep flocks longer in production, provided egg shell quality can be maintained."

He noted that "the increase in hen-housed egg production is mainly due to longer clutch length and improved uniformity of layer flocks. No doubt, there is a trend in cage layers to longer production cycles. A popular commercial goal is 500 eggs in one cycle with no moult, although this has already been surpassed in many flocks. The modern layer is capable of laying 150 eggs per clutch."

Dr Leeson, however, stressed that "genetic progress and longer laying cycles have consequences. Long laying cycle programmes start during pullet rearing – you can't make decisions at 72 weeks of age. Instead, you must start with your end goals, such as persistency, egg size and shell quality, in mind. You

can then develop a life-cycle approach to feeding, lighting, nutrition, and general management." Important issues to manage include:

Body weight control - early and late

Mature body weight dictates subsequent egg size. In the past, the common goal was being at, or above, management guide weight recommendations. For extended lay, a larger body weight results in too large an egg past 70 weeks of age, and so it is more difficult to maintain egg shell quality. Now the goal is to grow a slightly smaller pullet, and emphasis changes to achieving adequate early egg size from this smaller bird. This makes pre-lay nutrition for these slightly smaller pullets even more important.

The scheduling of rearing diets is more important than diet formulation. Dr Leeson's guidelines are:

- Starter diet 19-20% CP, 2,850-2,900 kcal ME/kg from day old to target pullet body weight
- Grower diet 17-18% CP, 2,800-2,900 kcal ME/kg from target body weight to mature body size
- Pre-lay diet (or layer diet?) 16-18% CP, 2,800-2,900 ME/kg, mature body size to first egg

All nutrients are important, but energy is usually limiting for egg number, whereas protein/amino acids influence egg size (and feathering).

There is now even more emphasis on pullet growing to ensure adequate fat reserves through peak production, so birds are in a positive energy balance. The establishment of an energy reserve occurs during the rearing phase and has a significant effect on the bird's body composition at point of lay.

Egg size control - early and late

The obvious solution to manage body weight (and egg size) is to light-stimulate a smaller pullet, or at least to not light-stimulate a heavy pullet. This achieves a balance between accepting reduced early egg size, versus limiting an increase in egg size late in the production cycle.

Egg size can be increased in smaller early-lay pullets by:

- Reducing environmental temperature, if possible, to stimulate feed intake
- Midnight feeding 19-29 weeks
- Adequate amino acid nutrition intake, tailored to feed intake, especially methionine
- Increased number of feedings/day and increased feed particle size (pellets)

Shell strength is negatively correlated with egg size. To temper egg size late in the cycle, Dr Leeson recommended:

- Body weight control
- Controlled day length: longer day length = increased feed intake, 14 hours maximum day length in controlled-environment houses
- Warmer temperature 26°C is ideal
- Reduce number of feedings and particle size
- Temper amino acid nutrition (with caution). Low crude protein/high amino acid diets limit the increase in egg size.

Midnight feeding provides about 1-hour extra light per day and therefore stimulating feed consumption in the middle of the dark period. Having access to feed during this period improves eggshell quality via the supply of calcium during the time when shell calcification takes place. The extra light period is perceived by the bird to be part of the night. The dark period after the light period must be longer than the initial dark period, as the bird perceives the start of the day is the end of the longest period of darkness. Removing midnight feeding should be done gradually – 15 minutes per week, advised Dr Leeson.

Preventing calcium depletion

Also known as cage layer fatigue, calcium depletion is becoming more common in all strains due to high sustained egg output. Calcium deficiency in the feed leads to loss of medullary or long bone (a reservoir of about 4g of calcium) and increased bone fragility. It is commonly seen at 35-40 weeks of age, with a 1-2% occurrence. If the incidence is more than 2%, seek advice for your pre-lay nutrition.

The development of the medullary bones takes about 10 days and requires additional calcium. Pre-lay rations support a smooth transition from developer feed to layer feed, with 2-2.5% calcium, while the other nutrients are similar to a layer feed. Pre-lay rations help the birds to adapt to the high calcium content of layer feed and to maintain sufficient daily feed intake.

To prevent calcium depletion, Dr Leeson suggested:

- Optimise pre-lay calcium (Ca) and phosphorous (P) nutrition
- Intake of 1.5g Ca, 350-450mg available P/day for at least 7 days prior to first egg
- During early lay, ensure 3.5-4 g Ca and 420 mg available P/day
- Consider vitamin D₃ water treatment (150 IU/day, twice weekly)

Pre-lay diets provide the bird with the opportunity to deposit medullary bone. This bone deposition coincides with follicular maturation and is under the control of both estrogens and androgens. The latter hormone seems essential for medullary bone growth, and its presence is manifested in the growth and reddening of the comb and wattles. Consequently, there will be little medullary deposition, regardless of diet calcium level, if the birds are not showing comb and wattle development and this stage of maturity should be the cue for increasing the bird's calcium intake.

Liver health

Excess energy relative to needs results in excess fat accumulation that is prone to oxidation. This is why you never see fatty liver haemorrhagic syndrome (FLHS) in poor-producing flocks. Layers normally have a very fatty liver, as 100% of egg yolk synthesis occurs in the liver.

The lower the fat content of the diet, the greater the stress/need to fat synthesis in the liver. With a low energy/low fat/carbohydrate diet FLHS is almost universal to varying degrees. One treatment is to add fat to the diet! Haemorrhage (not always FLHS) is inevitable with dietary omega-3s that are very prone to oxidation.

Dr Leeson recommended prevention/control for FLHS, which usually starts about weeks 36-40, including:

- +1.0 kg choline
- +0.5 kg methionine
- +100 IU vitamin E
- +30% does Hy-D because of impaired liver metabolism of vitamin D₃ (that can also impact calcium absorption)
- Add 2% dietary fat without change in diet energy level

<u>EW Nutrition</u>'s Poultry Academy took place in Jakarta and Manila in early September 2023. Dr. Steve Leeson, an expert in Poultry Nutrition & Production with nearly 50 years' experience in the industry, was the distinguished keynote speaker.

Dr. Leeson had his Ph.D. in Poultry Nutrition in 1974 from the University of Nottingham. Over a span of 38 years, he was a Professor in the Department of Animal &Poultry Science at the University of Guelph, Canada. Since 2014, he has been Professor Emeritus at the same University. As an eminent author, he has more than 400 papers in refereed journals and 6 books on various aspects of Poultry Nutrition & Management. He also won the American Feed Manufacturer's Association Nutrition Research Award (1981), the Canadian Society of Animal Science Fellowship Award (2001), and Novus Lifetime Achievement Award in Poultry Nutrition (2011).

Nutritional considerations for immunity and gut health



Conference report

At the recent EW Nutrition Poultry Academy in Jakarta, Indonesia, Dr Steve Leeson, Professor Emeritus, University of Guelph, Canada, opened his presentation by stating that "it is obvious that any nutrient deficiency will impact bird health, but not so obvious is that nutrition *per se* can positively impact immunity and health in an otherwise healthy and high-producing bird."

Modern high-performing broilers are characterized by extremely high feed intake. This puts a lot of stress on the physiology of the entire gastrointestinal tract, but particularly so on the absorptive epithelial cells of the small intestine. Any organism requires a nutrient source for survival and reproduction. Dr Leeson asked "can we significantly reduce nutrient supply to pathogens, while sustaining bird productivity?"

He reminded the audience that no cellular function comes for free: so there is always a "cost". A general

conclusion is that 10% of nutrients can be used for immune function during disease challenge, and always get priority. Therefore, you don't want to overstimulate the immune system, which in extreme situations leads to an inflammatory response. In his presentation, Dr Leeson considered factors determining gut health and nutritional tools which are available to support gut health.

Gut microflora

Gut pathogens impact the bird and/or the consumer. *Clostridia* and *E. coli* are the major concerns regarding bird health and productivity, whereas *Salmonella* and *Campylobacter* are major pathogens important for human health.

The chick hatches with a gut virtually devoid of microbes, so early colonizers tend to predominate quite quickly. Microbial species present on the hatching tray, during delivery and during the first few days at the farm will likely dictate early gut colonization. In some instances, the chick's microflora may be established by the time it gets to the farm, so the probiotic faces more of a challenge to establish itself as the predominant species.

Antibiotic alternatives

Gut villi development matures at around 10-15 days of age. The broiler pre-starter diet therefore is a target for feed additives that positively impact gut structure and development.

- Among the short chain fatty acids, butyric acid is considered the prime energy source for enterocytes and it is also necessary for the correct development of the gut-associated lymphoid tissue (GALT). Butyric acid can also be added indirectly via fermentation of judicious levels of soluble fiber to encourage optimal gut villi development. Dr Leeson added that he is a big believer in butyric acid, encouraging a good gut structure at 10 days, which can be worth about 50 kcal.
- **Exogenous enzymes** should also be considered in an attempt to maximize digestion and limit the flow of nutrients to the large intestine and ceca. Protease enzymes have great potential in this regard, since they allow nutritionists to reduce dietary crude protein and hopefully reduce the supply of nitrogen that fuels proteolytic Clostridia bacteria in the large intestine and ceca.
- Amino acids, particularly threonine, play a critical role in the maintenance of intestinal mucosal integrity and barrier function, especially for mucin synthesis, which protects enterocytes from adherence by pathogenic bacteria, and from attack by endogenous enzymes and acids.
- Polyunsaturated fatty acids (PUFAs) Omega-3s and especially DHA from fish oil help to reduce inflammatory response (overstimulation). Omega-3s are poorly converted to DHA by the chicken, so conventional sources such as flax are of limited application for immunity.
- Blood plasma from pigs or cattle is a complex spray-dried mixture of proteins and amino acids, many of which are immunoglobulins that "temper" the immune system, much like PUFAs.
- Vitamins A, D, E and C have vital roles in the normal function of the immune system and have antioxidant capacity.
- Certain complex carbohydrates, such as ß-glucans, influence gut health due to their fermentation, leading to the production of short-chain fatty acids, such as butyrate.
- Antioxidants to firstly control oxidation of fats and fat-soluble vitamins in feed, and secondly to optimize birds' cellular oxidative capacity, to prevent cell damage, therefore maintaining healthy cellular and immune function.
- Betaine increases intracellular water retention, reducing "dehydration" of microvilli and increasing their volume/surface area.
- **Fiber** moderate levels (1-2%) of soluble (fermentable) and insoluble fiber can be beneficial to early gut development by stimulating gizzard development and endogenous enzyme production.
- Phytogenics are becoming very common in combination with acidifiers (upper tract) and probiotics. Essential oils are becoming more mainstream the more we know about them.

Recommendations for optimizing gut health and immunity

Fast growth rate and high egg output are negatively correlated with immune response. Consequently, nutrient-dense diets are not optimal for immunity. With bacteria, it's a numbers game – but these numbers quickly multiply. The first 7 days are important, therefore probiotics must be established early. Consider the role of targeted feed additives, such as butyrate, phytogenics, antioxidants, PUFAs etc.

Also, maximize feed particle size – the limit is usually pellet quality. Mitigate nutrient transition at any diet change. Review the supply of trace minerals, as there is a trend to lower levels of organic minerals. With all the factors that weigh into production performance, any support that can be rallied through nutrition needs to be considered.

<u>EW Nutrition</u>'s Poultry Academy took place in Jakarta and Manila in early September 2023. Dr. Steve Leeson, an expert in Poultry Nutrition & Production with nearly 50 years' experience in the industry, was the distinguished keynote speaker.

Dr. Leeson had his Ph.D. in Poultry Nutrition in 1974 from the University of Nottingham. Over a span of 38 years, he was a Professor in the Department of Animal &Poultry Science at the University of Guelph, Canada. Since 2014, he has been Professor Emeritus at the same University. As an eminent author, he has more than 400 papers in refereed journals and 6 books on various aspects of Poultry Nutrition & Management. He also won the American Feed Manufacturer's Association Nutrition Research Award (1981), the Canadian Society of Animal Science Fellowship Award (2001), and Novus Lifetime Achievement Award in Poultry Nutrition (2011).

Meat labels explained



Certified Organic: (US, others) To be labeled as "Certified Organic" in the US, meat and poultry must come from animals that are raised in accordance with organic farming standards. These standards typically include restrictions on the use of synthetic pesticides, herbicides, antibiotics, and genetically modified organisms (GMOs). The animals are typically raised with organic feed and have access to the outdoors.

Chemical-free: (US) A product that contains no artificial ingredients or chemical preservatives.

Free-range or Free-roaming: (International) Poultry that has been allowed access to the outside.

Free-Range or Pasture-Raised: (US, others) These terms suggest that the animals had access to the outdoors or were raised on pasture, which can offer better living conditions than confined, industrial operations.

Fresh poultry: (US) Poultry that has never been below 26°F.

Frozen poultry: (US) Poultry that has been held at 0°F or lower.

Grain-Fed: (International) This label implies that the animals were primarily fed grains or other non-grassbased diets, which is common in many commercial meat production systems. **Grass-Fed**: (International) "Grass-Fed" typically means that the animals were primarily fed a diet of grass or forage throughout their lives, although some supplemental grains may be allowed. This label does not necessarily imply organic or non-GMO practices.

Halal: (International) Halal meat is prepared following Islamic dietary laws. This includes specific slaughter methods and requirements for the handling and preparation of the meat.

Kosher: (International) Kosher meat is prepared according to Jewish dietary laws and involves specific slaughtering practices and inspections.

Mechanically separated meat: (US) A paste-like meat product produced by forcing bones, with attached edible meat, under high pressure through a sieve or similar device to separate the bone from the edible meat tissue.

Natural: (US) A product containing no artificial ingredient or added color and is only minimally processed (a process which does not fundamentally alter the raw product). The label must explain the use of the term natural (such as "no added colorings or artificial ingredients; minimally processed").

No antibiotics: (US) The terms "no antibiotics added" may be used on labels for meat or poultry products if sufficient documentation is provided by the producer to the USDA demonstrating that the animals were raised without antibiotics. If an animal becomes sick and requires antibiotics, it cannot be sold as "no antibiotics added."

No hormones (beef): (US) The term "no hormones administered" may be approved for use on the label of beef products if sufficient documentation is provided to USDA by the producer showing no hormones have been used in raising the animals.

No hormones (pork or poultry): (US) Federal regulations prohibit the use of hormones in raising hogs and poultry.

Non-GMO: (International) A "Non-GMO" label indicates that the animals were not fed genetically modified organisms. This label may apply to both feed and the animals themselves.

Organic: (International) Meat and poultry labeled as organic must come from animals fed organic – which also means non-GMO – feed, given fresh air and outdoor access, and raised without antibiotics or added growth hormones. Organic livestock must also have access to pasture for at least 120 days per year.

Protected Designation of Origin (PDO) and Protected Geographical Indication (PGI): (EU) These labels are used to protect and promote regional and traditional foods. Meat labeled with PDO and PGI must come from specific regions and meet particular quality and production standards.

Raised without Antibiotics or Antibiotic-Free: (International) This label indicates that the animals were not treated with antibiotics during their lifetime. However, this label does not necessarily mean the animals were raised in organic or free-range conditions.

Sustainably Sourced: (International) This label may indicate that the meat was produced with a focus on environmental and ethical considerations, such as minimizing ecological impact and promoting fair labor practices.